


## NEUROLOGY IN PRACTICE

Series editors **Robert A. Gross & Jonathan W. Mink**

# Headache



Edited by

**Matthew S. Robbins, Brian M. Grosberg  
and Richard B. Lipton**

**WILEY** Blackwell



# Headache



NEUROLOGY IN PRACTICE:

SERIES EDITORS:

ROBERT A. GROSS, DEPARTMENT OF NEUROLOGY, UNIVERSITY OF ROCHESTER MEDICAL CENTER, ROCHESTER, NY, USA

JONATHAN W. MINK, DEPARTMENT OF NEUROLOGY, UNIVERSITY OF ROCHESTER MEDICAL CENTER, ROCHESTER, NY, USA

---

# Headache

EDITED BY

Matthew S. Robbins, MD

Brian M. Grosberg, MD

Richard B. Lipton, MD

*Montefiore Headache Center*

*Saul R. Korey Department of Neurology*

*Albert Einstein College of Medicine*

*Bronx, NY, USA*

**WILEY** Blackwell

This edition first published 2013, © 2013 by John Wiley & Sons, Ltd

*Registered office:* John Wiley & Sons, Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

*Editorial offices:* 9600 Garsington Road, Oxford, OX4 2DQ, UK  
The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK  
111 River Street, Hoboken, NJ 07030-5774, USA

For details of our global editorial offices, for customer services and for information about how to apply for permission to reuse the copyright material in this book please see our website at [www.wiley.com/wiley-blackwell](http://www.wiley.com/wiley-blackwell)

The right of the author to be identified as the author of this work has been asserted in accordance with the UK Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book. It is sold on the understanding that the publisher is not engaged in rendering professional services. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting a specific method, diagnosis, or treatment by health science practitioners for any particular patient. The publisher and the author make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of fitness for a particular purpose. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. Readers should consult with a specialist where appropriate. The fact that an organization or Website is referred to in this work as a citation and/or a potential source of further information does not mean that the author or the publisher endorses the information the organization or Website may provide or recommendations it may make. Further, readers should be aware that Internet Websites listed in this work may have changed or disappeared between when this work was written and when it is read. No warranty may be created or extended by any promotional statements for this work. Neither the publisher nor the author shall be liable for any damages arising herefrom.

ISBN: 9780470654729

A catalogue record for this book is available from the Library of Congress and the British Library.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Cover image: © iStockphoto.com/Sebastian Kaulitzki  
Cover design by Sarah Dickinson

Set in 8.75/11.75pt Utopia by Toppan Best-set Premedia Limited, Hong Kong

# Contents

|                  |            |
|------------------|------------|
| Contributor List | <i>vii</i> |
| Series Foreword  | <i>x</i>   |
| Preface          | <i>xi</i>  |
| Acknowledgments  | <i>xii</i> |

## PART I: PRINCIPLES OF HEADACHE: PRIMARY AND SECONDARY HEADACHE DISORDERS

|  |           |
|--|-----------|
| <b>1 The Basics of Headache Classification and Diagnosis</b>           | <b>3</b>  |
| <i>Seymour Solomon and Richard B. Lipton</i>                           |           |
| <b>2 Approach to the Patient with Headache</b>                         | <b>16</b> |
| <i>Brian M. Grosberg, Benjamin W. Friedman, and Seymour Solomon</i>    |           |
| <b>3 Secondary Headache Disorders Encountered in Clinical Practice</b> | <b>27</b> |
| <i>Deborah I. Friedman</i>   |           |

## PART II: MIGRAINE

|  |            |
|--|------------|
| <b>4 Diagnosis and Subtypes of Migraine</b>                                      | <b>57</b>  |
| <i>Seymour Solomon and Brian M. Grosberg</i>                                     |            |
| <b>5 Epidemiology, Progression, Prognosis, and Comorbidity of Migraine</b>       | <b>64</b>  |
| <i>Richard B. Lipton, C. Mark Sollars, and Dawn C. Buse</i>                      |            |
| <b>6 Pathophysiology and Genetics of Migraine</b>                                | <b>88</b>  |
| <i>Peter J. Goadsby</i>  |            |
| <b>7 Multidisciplinary Approach to Patients with Migraine</b>                    | <b>100</b> |
| <i>Lucille A. Rathier, Dawn C. Buse, Robert A. Nicholson, and Frank Andrasik</i> |            |
| <b>8 Acute Treatments for Migraine</b>   | <b>113</b> |
| <i>Dagny Holle and Hans-Christoph Diener</i>                                     |            |
| <b>9 Preventive Treatments for Migraine</b>                                      | <b>125</b> |
| <i>William B. Young</i>  |            |
| <b>10 Managing the Special Problem of Chronic Migraine</b>                       | <b>140</b> |
| <i>Marcelo E. Bigal</i>  |            |

## PART III: TENSION-TYPE HEADACHE

|   |            |
|---|------------|
| <b>11 Diagnosis, Subtypes, Epidemiology, Progression, Prognosis, and Comorbidity of Tension-type Headache</b> | <b>155</b> |
| <i>Sara C. Crystal, Uri Napchan, and Matthew S. Robbins</i>   |            |

|   |     |
|---|-----|
| 12 Pathophysiology and Genetics of Tension-type Headache                                      | 164 |
| <i>Sait Ashina and Lars Bendtsen</i>  |     |
| 13 Treatment of Tension-type Headache   | 172 |
| <i>Sara C. Crystal and Katherine A. Henry</i>   |     |
| PART IV: TRIGEMINAL AUTONOMIC CEPHALALGIAS INCLUDING CLUSTER HEADACHE                         |     |
| 14 Diagnosis and Subtypes of Trigeminal Autonomic Cephalalgias                                | 183 |
| <i>Arne May</i>   |     |
| 15 Epidemiology, Progression, Prognosis, and Comorbidity of Trigeminal Autonomic Cephalalgias | 192 |
| <i>Matthew S. Robbins and Jessica Ailani</i>  |     |
| 16 Pathophysiology and Genetics of Trigeminal Autonomic Cephalalgias                          | 201 |
| <i>Peter J. Goadsby</i>   |     |
| 17 Treatment of Trigeminal Autonomic Cephalalgias Including Cluster Headache                  | 214 |
| <i>Sarah Vollbracht and Brian M. Grosberg</i>   |     |
| PART V: OTHER HEADACHE DISORDERS  |     |
| 18 New Daily Persistent Headache  | 229 |
| <i>Matthew S. Robbins</i>   |     |
| 19 Hemicrania Continua  | 236 |
| <i>Uri Napchan</i>  |     |
| 20 Unusual Short-duration Primary Headaches   | 245 |
| <i>Sarah Vollbracht and Brian M. Grosberg</i>   |     |
| PART VI: MANAGEMENT OF HEADACHE IN SPECIFIC PATIENT POPULATIONS                               |     |
| 21 Management of Headache in Women  | 259 |
| <i>Elizabeth W. Loder, Dawn C. Buse, Vince Martin, Luzma Cardona, and Dawn A. Marcus</i>      |     |
| 22 Management of Headache in Children   | 269 |
| <i>Oranee Sanmaneechai and Karen Ballaban-Gil</i>   |     |
| 23 Management of Headache in the Elderly  | 290 |
| <i>Matthew S. Robbins and Richard B. Lipton</i>   |     |
| Index   | 307 |



# Contributor List

**Jessica Ailani, MD**

Assistant Professor, Department of Neurology  
Director, Georgetown University Headache  
Center  
Georgetown University Hospital  
Washington, DC, USA

**Frank Andrasik, PhD**

Distinguished Professor and Chair  
Department of Psychology  
University of Memphis  
Memphis, TN, USA

**Sait Ashina**

Headache Program, Director  
Assistant Professor of Neurology and  
Anesthesiology  
Department of Pain Medicine and Palliative  
Care  
Department of Neurology  
Beth Israel Medical Center  
Albert Einstein College of Medicine  
New York, NY, USA

**Karen Ballaban-Gil, MD**

Professor of Clinical Neurology and Clinical  
Pediatrics  
Albert Einstein College of Medicine  
Montefiore Medical Center  
Bronx, NY, USA

**Lars Bendtsen, MD, PhD**

Associate Professor  
Danish Headache Center and Department of  
Neurology,  
University of Copenhagen, Glostrup Hospital,  
Glostrup, Copenhagen, Denmark

**Marcelo E. Bigal, MD, PhD**

Head of the Merck Investigator Studies Program,  
Scientific Engagements and Education (MISP/  
SEE)  
North Wales, PA, USA, and  
Clinical Associate Professor, The Saul R. Korey  
Department of Neurology, Albert Einstein  
College of Medicine  
Bronx, NY, USA

**Dawn C. Buse, PhD**

Associate Professor, Department of Neurology  
Albert Einstein College of Medicine of Yeshiva  
University,  
Assistant Professor, Clinical Health Psychology  
Doctoral Program  
Ferkauf Graduate School of Psychology of  
Yeshiva University  
Bronx, NY, and  
Director of Behavioral Medicine  
Montefiore Headache Center  
Bronx, NY, USA

**Luzma Cardona, MD**

Instructor in Neurology  
Harvard Medical School  
Boston, MA, and  
Staff Physician  
Graham Headache Centre  
Faulkner Hospital  
Boston, MA, USA

**Sara C. Crystal, MD**

Clinical Assistant Professor  
Department of Neurology  
NYU School of Medicine  
New York Headache Center  
New York, NY, USA

**Hans-Christoph Diener, MD, PhD**

Department of Neurology and Headache Center  
University Duisburg-Essen  
Essen, Germany

**Benjamin W. Friedman, MD**

Attending Physician, Department of Emergency  
Medicine  
Montefiore Medical Center, Bronx, NY, and  
Associate Professor of Emergency Medicine  
Albert Einstein College of Medicine  
Bronx, NY, USA

**Deborah I. Friedman, MD, MPH**

Professor, Department of Neurology &  
Neurotherapeutics and Ophthalmology  
University of Texas Southwestern  
Dallas, TX, USA

**Peter J. Goadsby, MD, PhD**

Headache Group, Department of Neurology  
University of California, San Francisco  
UCSF Headache Center  
San Francisco, CA, USA

**Brian M. Grosberg, MD**

Associate Professor of Neurology  
Albert Einstein College of Medicine, and  
Co-Director, Montefiore Headache Center  
Director, Headache and Facial Pain Fellowship  
Montefiore Headache Center  
Bronx, NY, USA

**Katherine A. Henry, MD**

Associate Professor  
Department of Neurology  
New York University School of Medicine  
New York, NY, USA

**Dagny Holle, MD**

Department of Neurology and Headache Center  
University Duisburg-Essen  
Essen, Germany

**Richard B. Lipton, MD**

Edwin S. Lowe Professor and Vice Chair of  
Neurology  
Professor of Epidemiology and Population  
Health,  
Professor of Psychiatry and Behavioral Science  
Albert Einstein College of Medicine, and  
Co-Director, Montefiore Headache Center  
Bronx, NY, USA

**Elizabeth W. Loder, MD, MPH**

Associate Professor of Neurology  
Harvard Medical School  
Boston, MA, USA, and  
Chief, Division of Headache and Pain  
Department of Neurology  
Brigham and Women's/Faulkner Hospitals  
Boston, MA, USA

**Dawn A. Marcus, MD**

Professor, Department of Anesthesiology  
University of Pittsburgh  
Pittsburgh, PA, and  
Headache Clinic  
University of Pittsburgh  
Pittsburgh, PA, USA

**Vince Martin, MD**

Professor, Department of Internal Medicine,  
University of Cincinnati  
and  
Headache Clinic,  
University of Cincinnati  
Cincinnati, OH, USA

**Arne May, MD**

Professor of Neurology  
Department of Systems Neuroscience  
Universitäts-Krankenhaus Eppendorf (UKE)  
Hamburg, Germany

**Uri Napchan, MD**

Middletown Medical  
Middletown, NY, USA

**Robert A. Nicholson, PhD**

Associate Professor of Research  
Department of Neurology & Psychiatry and  
School of Public Health  
Saint Louis University School of Medicine, and  
Senior Research Methodology Advisor  
Mercy Health Research & Ryan Headache  
Center  
St. Louis, MO, USA

**Lucille A. Rathier, PhD**

Clinical Assistant Professor  
Psychiatry and Human Behavior  
Warren Alpert Medical School, Brown  
University, and  
Staff Psychologist  
Behavioral Medicine Clinical Services  
The Miriam Hospital  
Providence, RI, USA

**Matthew S. Robbins, MD**

Assistant Professor of Neurology  
Albert Einstein College of Medicine  
Chief of Service—Neurology  
Einstein Division of Montefiore Medical Center,  
and  
Director of Inpatient Services  
Montefiore Headache Center  
Bronx, NY, USA

**Oranee Sanmaneechai, MD**

Fellow in Child Neurology  
Albert Einstein College of Medicine, and  
Assistant Professor of Pediatrics  
Division of Neurology, Department of Pediatrics  
Siriraj Hospital, Mahidol University  
Bangkok, Thailand

**C. Mark Sollars, MS**

Research Coordinator  
Montefiore Headache Center  
Bronx, NY, USA

**Seymour Solomon, MD**

Director Emeritus, Montefiore Headache Center  
Montefiore Medical Center, Bronx, NY, and  
Professor of Neurology  
Albert Einstein College of Medicine  
Bronx, NY, USA

**Sarah Vollbracht, MD**

Assistant Professor of Neurology  
Albert Einstein College of Medicine  
Montefiore Headache Center  
Bronx, NY, USA

**William B. Young, MD, FAHS, FAAN**

Professor of Neurology  
Thomas Jefferson University  
Philadelphia, PA, USA

# Series Foreword

The genesis for this book series started with the proposition that, increasingly, physicians want direct, useful information to help them in clinical care. Textbooks, while comprehensive, are useful primarily as detailed reference works but pose challenges for uses at the point of care. By contrast, more outline-type references often leave out the “hows and whys”—pathophysiology, pharmacology—that form the basis of management decisions. Our goal for this series is to present books, covering most areas of neurology, that provide enough background information to allow the reader to feel comfortable, but not so much as to be overwhelming; and to associate that with practical advice from experts about care, combining the growing evidence base with best practices.

Our series will encompass various aspects of neurology, with topics and the specific content chosen to be accessible and useful.

Chapters cover critical information that will inform the reader of the disease processes and mechanisms as a prelude to treatment planning. Algorithms and guidelines are presented, when appropriate. “Tips and Tricks” boxes provide expert suggestions, while other boxes present cautions and warnings to avoid pitfalls. Finally, we provide “Science Revisited” sections that review the most important and relevant science background material, and “Bibliography”

sections that guide the reader to additional material.

We welcome feedback. As additional volumes are added to the series, we hope to refine the content and format so that our readers will be best served.

Our thanks, appreciation, and respect go out to our editors and their contributors, who conceived and refined the content for each volume, assuring a high-quality, practical approach to neurological conditions and their treatment.

Our thanks also go to our mentors and students (past, present, and future), who have challenged and delighted us; to our book editors and their contributors, who were willing to take on additional work for an educational goal; and to our publisher, Martin Sugden, for his ideas and support, for wonderful discussions and commiseration over baseball and soccer teams that might not quite have lived up to expectations. We would like to dedicate the series to Marsha, Jake, and Dan; and to Janet, Laura, and David. And also to Steven R. Schwid, MD, our friend and colleague, whose ideas helped to shape this project and whose humor brightened our lives, although he could not complete this goal with us.

Robert A. Gross  
Jonathan W. Mink  
*Rochester, NY, USA*

# Preface

The practice of clinical neurology has changed remarkably over the last 25 years, transforming from a diagnosis-oriented specialty into a discipline with many dynamic therapeutic opportunities. The subspecialty of headache medicine exemplifies this transformation as insights from genetics, imaging, and animal models have supported the development of specific treatments for migraine and other headache disorders. As a consequence, clinicians have an unparalleled opportunity to relieve pain, restore function, and improve quality of life for our patients.

Headache is such an extraordinarily common complaint encountered in medical settings, having the unusual duality of often being both intimidating and disregarded. In some patients, headaches may be an infrequent nuisance, which responds well to over-the-counter medications. In others, headaches may occur on a daily basis and completely disrupt educational, occupational, and social function, despite treatment. Diagnosis is complicated by the hundreds of different etiologies, both benign and life-threatening, that give rise to headache.

In this volume, we present a practical overview of headache disorders, designed to address the needs of both neurologists and general practitioners alike. Our authors include both research experts from the laboratory and practitioners who are “in the trenches.” In the first section, we review the principles of headache medicine by focusing on the classification and diagnostic approach of both primary and secondary headache disorders.

The second section features a comprehensive, multifaceted appraisal of migraine, the most common headache disorder that presents to medical attention. Tension-type headache is the focus of the third section, a disorder that is usually quite benign although enigmatic, but in some patients can require significant multidisciplinary treatment. We then present the trigeminal autonomic cephalalgias, which while uncommon are the most distinctive group of headache disorders and require some expertise in diagnosis and treatment.

In the fifth section, we review uncommon and unusual primary headache disorders, including new daily persistent headache and hemicrania continua. We conclude by detailing the approach to managing headache disorders in specific populations, including women, children, and the elderly.

Numerous volumes addressing headache have been published previously. The goal of this work is to present a crisp, practical approach to headache disorders that has been written by a diverse group of experienced clinicians and scientists. We hope our approach resonates with both trainees, newly credentialed practitioners, and experienced clinicians in the many medical fields where headache is encountered.

Matthew S. Robbins, MD  
Brian M. Grosberg, MD  
Richard B. Lipton, MD

# Acknowledgments

The successes and accomplishments of the authors would not have been possible without the support and sacrifices of many other people. All three authors would like to thank their colleagues at the Montefiore Headache Center: Seymour Solomon, Dawn Buse, Sarah Vollbracht, Jelena Pavlovic and Mark Sollars for their friendship and brilliance.

Matthew Robbins thanks his lovely wife Hilary for her supreme devotion, partnership, support, wisdom, and guidance in all aspects of life, and for sharing in the joy of raising their magnificent children Zoe and Dylan. His parents Mitchell and Debra, brother David, and grandmother Rose have remarkably provided an unparalleled lifetime of love and cheerleading. His co-editors of this volume have been two of his major professional mentors and he expresses eternal gratitude towards them.

Brian Grosberg would like to thank his mentors Seymour Solomon, Richard Lipton, and Deborah Friedman for teaching him with untiring enthusiasm and for encouraging him to be the best he can be. He is amazingly grateful to his family: his parents Carole and Lee Grosberg and his brother Michael for their unending love and support, his

children Noah and Talia for the joy they bring him every moment of every day and who have been waiting patiently for Daddy to finish this book and be able to see their names in print, and most of all he acknowledges the love, support, companionship, sacrifice and encouragement of his wife Jennifer who is the love of his life and makes everything he does possible.

Richard Lipton thanks Amy, his wife and partner for 36 years, for being his compass, for giving him deep roots, wonderful children and so much love. He thanks his children, Lianna and Justin, for the joyful chapters lived together and for all the promise contained in their every smile. He thanks his research mentor, Philip Holzman, for teaching him what it means to be a scientist and his clinical mentor, Seymour Solomon, for teaching him the ways and pleasures of relieving pain. He also thanks his many collaborators in headache medicine, particularly Walter Stewart, Marcelo Bigal, Dawn Buse and Daniel Serrano for teaching him the joys of shared discovery. Finally, he thanks his co-editors of this volume for being inspirational clinicians and teachers and for their infinite patience and hard work in bringing this book to fruition.

Part I

# Principles of Headache: Primary and Secondary Headache Disorders





# The Basics of Headache Classification and Diagnosis

Seymour Solomon<sup>1,2</sup> and Richard B. Lipton<sup>1,2</sup>

<sup>1</sup>Montefiore Headache Center, Bronx, NY, USA

<sup>2</sup>Albert Einstein College of Medicine, Bronx, NY, USA

## Introduction

The causes of headache range from the short-lived and trivial (e.g., a hangover headache) to the intermittent and quality-of-life threatening (migraine), to the unremitting and life-threatening (subarachnoid hemorrhage headache). This broad range of etiologies is matched by the very high prevalence of headache in the general population. In combination, the diversity of causes and high prevalence mandate a systematic approach to classification and diagnosis. Classification refers to a set of categories with diagnostic rules that provide the framework for a clinical approach. Diagnosis is the process of applying the rules to individual patients, defining their place in the classification.

In this chapter, we present an approach to both headache classification and diagnosis. We begin by describing the classification for headache disorders (see Tables 1.1, 1.2, 1.3, and 1.4 in the relevant sections). We recommend a three-step diagnostic process. First, we emphasize the identification or exclusion of secondary headache disorders by history, physical examination, and judicious use of diagnostic tests (see Table 1.5 and Figure 1.1). Second, we consider four groups of primary headache disorders that are defined

based on headache frequency and duration (see Table 1.6) and refer to these as primary headache syndromes. Finally, we emphasize the identification of specific disorders within syndromic groups.

## Approach to classification

The classification system for headache disorders has evolved over the past 50 years. Good classification systems should be valid, reliable, generalizable, and complete. A valid system provides categories that correspond, as much as possible, to biologic reality. As a practical matter, a valid system should usefully predict prognosis, response to treatment, and pathobiology. In a reliable system, two clinicians seeing the same patient should assign the same diagnoses. A generalizable system should work in a variety of settings including population studies, primary care, specialty care, and clinical trials. Finally, a complete system has an appropriate category for every headache type.

The first modern attempt at headache classification, sponsored by the National Institute of Health in 1962, was undertaken by the Ad Hoc Committee on Headache Classification of Headache [1]. While the Ad Hoc Committee's

descriptions of headache types, for the most part, stand to this day, disorders were described in terms of their typical features without explicit operational rules regarding the characteristics required to diagnose or exclude particular disorders. In the absence of such rules, diagnostic reliability is not possible.

For the past 25 years, headache classification efforts have been led by the International Headache Society, which has published two editions of its International Classification of Headache Dis-

orders (known as ICHD-1 and -2) [2, 3]. A third edition (ICHD-3) is in development. Learning the lessons taught by the Diagnostic and Statistical Manual of Mental Disorders system in psychiatry, the ICHD system provides clear boundaries among the primary headache disorders (Table 1.1). Although it is based on both evidence and expert opinion, the effort to be explicit inevitably leads to rules that are somewhat arbitrary. For example, to diagnose migraine without aura, at least five headache attacks are required. A patient

**Table 1.1.** The classification system (modified) (numbers refer to the ICHD-2 code)

### **A. Primary headache disorders**

1. Migraine
  - 1.1. Migraine without aura
  - 1.2. Migraine with aura
  - 1.3. Childhood periodic syndromes that are commonly precursors with migraine
  - 1.4. Retinal migraine
  - 1.5. Complications of migraine
  - 1.6. Probable migraine
2. Tension-type headache
  - 2.1. Infrequent episodic tension-type headache
  - 2.2. Frequent tension-type headache
  - 2.3. Chronic tension-type headache
  - 2.4. Probable tension-type headache
3. Cluster headache and other trigeminal autonomic cephalalgias (TAC)
  - 3.1. Cluster headache
    - 3.1.1. Episodic cluster headache
    - 3.1.2. Chronic cluster headache
  - 3.2. Paroxysmal hemicrania
    - 3.2.1. Episodic paroxysmal hemicrania
    - 3.2.2. Chronic paroxysmal hemicrania
  - 3.3. Short-lasting unilateral neuralgiform headache attacks with conjunctival injection and tearing (SUNCT)
  - 3.4. Probable TAC
4. Other primary headaches
  - 4.1. Primary stabbing headache
  - 4.2. Primary cough headache
  - 4.3. Primary exertional headache
  - 4.4. Primary headache associated with sexual activity
  - 4.5. Hypnic Headache
  - 4.6. Primary thunderclap headache
  - 4.7. Hemicrania continua
  - 4.8. New daily persistent headache (NDPH)

### **B. Secondary headache disorders**

5. Headache attributed to head and/or trauma
  - 5.1. Acute post-traumatic headache
  - 5.2. Chronic post-traumatic headache

**Table 1.1.** (Continued)

6. Headache attributed to cranial or cervical vascular disorder
  - 6.1. Headache attributed to ischemic stroke or transient ischemic attack
  - 6.2. Headache attributed to non-traumatic intracranial hemorrhage
  - 6.3. Headache attributed to unruptured vascular malformation
  - 6.4. Headache attributed to arteritis
  - 6.5. Carotid or vertebral artery pain
  - 6.6. Headache attributed to cerebral venous thrombosis
  - 6.7. Headache attributed to other intracranial vascular disorder
7. Headache attributed to non-vascular intracranial disorder
  - 7.1. Headache attributed to high cerebrospinal fluid pressure
  - 7.2. Headache attributed to low cerebrospinal fluid pressure
  - 7.3. Headache attributed to non-infectious inflammatory disease
  - 7.4. Headache attributed to intracranial neoplasm
  - 7.5. Headache attributed to intrathecal injection
  - 7.6. Headache attributed to epileptic seizure
  - 7.7. Headache attributed to Chiari malformation type I
  - 7.8. Syndrome of transient headache and neurologic deficits with cerebrospinal fluid lymphocytosis
  - 7.9. Headache attributed to other non-vascular intracranial disorder
8. Headache attributed to a substance or its withdrawal
  - 8.1. Headache induced by acute substance use or exposure
  - 8.2. Medication-overuse headache (MOH)
  - 8.3. Headache as an adverse event attributed to chronic medication
  - 8.4. Headache attributed to substance withdrawal
9. Headache attributed to infection
  - 9.1. Headache attributed to intracranial infection
  - 9.2. Headache attributed to systemic infection
  - 9.3. Headache attributed to HIV/AIDS
  - 9.4. Chronic post-infection headache
10. Headache attributed to disorder of homeostasis
  - 10.1. Headache attributed to hypoxia and/or hypercapnia
  - 10.2. Dialysis Headache
  - 10.3. Headache attributed to hypertension
  - 10.4. Headache attributed to hypothyroidism
  - 10.5. Headache attributed to fasting
  - 10.6. Cardiac cephalgia
  - 10.7. Headache attributed to other disorder of homeostasis

**C. Headache or facial pain attributed to disorder of cranial structures, psychiatric disorders, cranial neuralgias**

11. Headache or facial pain attributed to disorder of cranium, neck, eyes, ears, nose, sinuses, teeth, mouth or other facial or cranial structures
  - 11.1. Headache attributed to disorder of cranial bone
  - 11.2. Headache attributed to disorder of neck
  - 11.3. Headache attributed to disorder of eyes
  - 11.4. Headache attributed to disorder of ears
  - 11.5. Headache attributed to rhinosinusitis
  - 11.6. Headache attributed to disorder of teeth, jaws or related structures
  - 11.7. Headache or facial pain attributed to temporomandibular joint disorder
  - 11.8. Headache attributed to other disorder of cranium, neck, eyes, ears, nose sinuses, teeth, mouth or other facial or cervical structures

(Continued)

**Table 1.1.** (Continued)

12. Headache attributed to psychiatric disorder
  - 12.1. Headache attributed to somatisation disorder
  - 12.2. Headache attributed to psychotic disorder
13. Cranial neuralgias and central causes of facial pain
  - 13.1. Trigeminal neuralgia
  - 13.2. Glossopharyngeal neuralgia
  - 13.3. Nervus intermedius neuralgia
  - 13.4. Superior laryngeal neuralgia
  - 13.5. Nasociliary neuralgia
  - 13.6. Supraorbital neuralgia
  - 13.7. Other terminal branch neuralgias
  - 13.8. Occipital neuralgia
  - 13.9. Neck-tongue syndrome
  - 13.10. External compression headache
  - 13.11. Cold-stimulus headache
  - 13.12. Constant pain caused by compression, irritation or distortion of cranial nerves or upper cervical roots by structural lesions
  - 13.13. Optic neuritis
  - 13.14. Ocular diabetic neuropathy
  - 13.15. Head or facial pain attributed to herpes zoster
  - 13.16. Tolosa-Hunt syndrome
  - 13.17. Ophthalmoplegic “migraine”
  - 13.18. Central causes of facial pain
  - 13.19. Other cranial neuralgia or other centrally mediated facial pain
14. Other headache, cranial neuralgia, central or primary facial pain
  - 14.1. Headache not elsewhere classified
  - 14.2. Headache unspecified

with four typical attacks of migraine and a family history of migraine almost certainly has migraine, although they may not meet full criteria for the disorder.

Since ICHD-1, the criteria have evolved, based on formal field testing and clinical experience [3]. Major improvements have been made in the area of completeness as important disorders not included in ICHD-1, such as chronic migraine, new daily persistent headache, and hypnic headache were added to ICHD-2, published in 2004 [3]. In the years that followed, as field testing revealed limitations to ICHD-2, proposed revisions have been published, which will culminate in ICHD-3. This chapter is based on ICHD-2 and the published proposals for revision that will inform the shape of ICHD-3.

The ICHD approach to classification defines three overarching categories: primary headache disorders, secondary headache disorders, and cranial neuralgias and facial pains. For the primary disorders, the headache disorder is the problem; there is no underlying disease. For secondary headaches, the headache is attributed to an underlying cause. In this chapter, the discussion of secondary headaches will deal mainly with the features that differentiate them from the primary headaches, i.e., the “red flags” that indicate the need for further diagnostic evaluation. The primary headaches are divided into four broad groups: migraine, tension-type headache (TTH), trigeminal autonomic cephalalgias (TACs; including cluster headache), and other uncommon primary headaches.

## Approach to diagnosis

### Headache history

Because most patients seeing their doctor for headache have normal medical and neurologic examinations, the history is essential for accurate diagnosis. We recommend beginning with an open-ended statement such as, “Tell me about your headaches and how they affect your life.” Although many clinicians fear that such an open-ended statement will generate a very long and unhelpful narrative, most patients speak for less than 90 seconds and usually provide useful information about both the headache’s features and the impact on their life. Studies show that the use of open-ended questions makes headache visits shorter and increases both clinician and patient satisfaction with the visit.

After the patient’s narrative, directed questions are used to fill in the details. The age of onset of the headache and the accompanying circumstances are important. Ask about the features of the head pain, including the location (unilateral, bilateral, or focal), quality (throbbing, stabbing, or steady ache), and intensity of the pain (on a scale of 1 to 10). The duration and frequency of headaches are important. The timing of attacks may assist both diagnosis and therapy. What time in the 24-hour day do they occur; are they associated with menstruation; do they cluster in time?

#### ★ TIPS AND TRICKS

Patients often first describe their worst or most recent headache. They may rate the intensity as 10 on a scale of 1 to 10, and the duration as constant since last Tuesday. Ask about the range of intensity and the typical duration. Similarly, they may state that the headache lasts for 3 days and occurs three times per week. It then is useful to determine the number of headache-free days per week or per month.

Ask about possible warning or premonitory symptoms that typically precede the onset of the headache by hours or days. Typical premonitory features include changes in mood such as irritability, food cravings, and neck stiffness among others. Auras are focal neurologic symptoms that

usually begin 15–60 minutes before the headache. Premonitory features and auras are most typical of migraine although not entirely specific. Symptoms accompanying the headache may include nausea, vomiting, photophobia, and phonophobia as well as tearing of the eye, conjunctival injection, and nasal discharge among others.


It is also helpful to understand factors associated with an increased chance of headache onset (triggers). Avoidance or management of triggers becomes a therapeutic strategy. Aggravating factors increase the severity of a pre-existing headache but are not associated with an increased rate of headache initiation. Ameliorating factors provide diagnostic clues. For example, hibernation in a dark, quiet room suggests a diagnosis of migraine. Understanding the effects of treatment and the course of the headache over time is also helpful.

#### ★ TIPS AND TRICKS

Patients often list their prescription medication, but fail to mention the over-the-counter (OTC) medications they use. OTC medications and dietary caffeine are important causes of medication overuse headache. After the history of the present illness has been obtained, other historical factors should be elicited. Is there a past history of illness, surgery, trauma, depression, or anxiety? Family history should include questions on other members with headache. Social history factors relating to marital status, education, occupation, interests, habits (alcohol, nicotine, and illicit drugs), sleep patterns, childhood abuse, and past and present stressors may all be relevant.

### Diagnosing headache is as easy as 1, 2, 3

The first step in evaluating a patient with headache is to differentiate primary from secondary headaches. Secondary headaches are attributable to an underlying neurologic, medical, or psychiatric illness (Figure 1.1). Secondary headache is suspected if red flags on the history or examination suggest it. Primary headache disorders are likely if red flags are absent or if diagnostic testing excludes secondary headache. In addition, if a



**Figure 1.1.** Headache diagnosis. (Reproduced from *Wolff's Headache and Other Pain*, 8th edition by Lipton et al. (2008) [15]. By permission of Oxford University Press, USA.)

secondary cause of headache is present but temporally remote from a pre-existing headache disorder, both primary and secondary headaches are likely. Consider a woman with typical attacks of migraine with aura from age 10 who develops a new form of daily headache at age 70. The MRI reveals a meningioma. Most likely, the patient has migraine with aura and a secondary headache attributed to her meningioma.

#### ★ TIPS AND TRICKS

When a headache changes in intensity or frequency, or in another feature, it may be a worsening of the primary headache, but one must always consider the superimposition of a new secondary headache. Searching for a new potential cause is warranted.

In making the diagnosis of a primary headache, the history and physical and neurologic examinations do not suggest secondary headache, or the latter can be ruled out either by investigations or by a distant temporal relationship (see Table 1.5). Symptoms that point to a specific primary headache but may not include all of the diagnostic criteria listed in ICHD-2 are preceded by the title “probable,” for example “probable migraine.” If the patient has two or more types of headache contemporaneously, all should be classified. The presence of a primary headache does not, of course, exclude the development of a secondary headache. Inattention to that obvious fact is a common and sometimes fatal error.

After excluding a secondary headache disorder, the next step is to identify a primary headache syndrome. Common headache syndromes and the specific disorders to consider within each syndromic group are summarized in Table 1.6.

These syndromes are defined based on the number of headache days per month and on the average duration of attacks. Based on headache days, two examples with long and short frequencies are chronic TTH, with 15 or more days per month, and episodic TTH with fewer than 15 headache days per month. Based on duration, we consider short-duration attacks as lasting less than 4 hours and long-duration attacks as lasting for 4 hours or more, for example cluster headache and chronic migraine, respectively. Some ICHD-2 defined disorders are included in more than one syndromic group. Most people with migraine have episodic headache of a long duration, but a clinically important subgroup have chronic migraine, the most important cause of chronic daily headache with a long attack duration.

Having identified the syndrome, the next task is to diagnose the specific disorder within the syndromic group. In the sections that follow, we discuss a number of specific primary headache disorders.

### **Migraine (ICHD-2 1.0)**

Migraine is divided into two categories based on the characteristics of the attack: migraine without aura and migraine with aura. Attacks typically occur from less than once to several times a month. Migraine is also categorized based on headache days per month. When migraine occurs on 15 or more days per month for at least 3 months, the term “chronic migraine” is applied. The term “chronic” has several possible meanings in ordinary English and in medicine. As applied to migraine and TTH, it has the very specific meaning of headaches occurring on more days than not over at least 3 months. There are rare variants of migraine: retinal migraine, familial or sporadic hemiplegic migraine, and basilar-type migraine. Migraine is a very common phenomenon, affecting 12% of the population—18% of women and 6% of men in any given year. In contrast to past concepts, epidemiologic studies have shown that the prevalence of migraine varies inversely with socioeconomic status [4].

#### **Migraine without aura**

Migraine without aura (ICHD-2 1.1) represents about 75%–80% of all migraineurs. The criteria for migraine without aura are listed in Table 1.2. The typical features of an attack are a headache

that is one-sided, pulsating, moderate or severe in pain intensity, lasting for hours, and associated with nausea and/or vomiting, photophobia, and phonophobia. In children, the duration is often shorter, 1 or 2 hours, and the associated features of nausea or vomiting or both are sometimes more prominent than the headache. Migraine attacks may not be typical but still fulfill the diagnostic criteria. For example, the location is often bilateral, and the quality may be a steady ache rather than pulsating. In the absence of these two features, the diagnosis is still appropriate if the two other features of the headache are present (pain of moderate or severe intensity and aggravation by physical activity) and other criteria are fulfilled. Similarly, only one of the two associated features—nausea and sensitivity to light and sound—need be present. The frequency of attacks may vary from one per week to one per year. When the frequency increases to more than 15 days per month, the additional classification of chronic migraine is applied.

#### **Migraine with aura**

Migraine with aura (ICHD-2 1.2) (Table 1.2) occurs in about 20% of people with migraine. The most common aura of migraine is a positive homonymous visual phenomenon, a scintillating arc of zigzag lines that occurs for minutes before the onset of the headache and then completely disappears. Any neurologic symptom or sign may be an aura of migraine, but unusual conditions are more specifically subclassified as hemiplegic migraine, basilar-type (brainstem dysfunction) migraine, or retinal migraine (see “Subtypes of migraine”). The symptoms are usually positive, such as flickering lights or tingling sensations, in contrast to negative symptoms such as loss of vision or impaired sensation. The visual symptoms are typically homonymous, and the sensory symptoms usually affect the hand and face. These focal symptoms are most often on the side opposite the headache. Dysphasic speech is the least common aura manifestation. The aura usually evolves gradually and lasts from 5 to 60 minutes. Headache with features of migraine without aura usually begins as soon as the aura stops but sometimes begins at the same time as the aura and may begin as long as 1 hour after the aura. The aura may occur without headache, especially during middle or late age. Typical auras

**Table 1.2.** Diagnostic criteria for migraine without and with aura**ICHD-2 diagnostic criteria for 1.1 migraine without aura**

- A. At least 5 attacks fulfilling criteria B–D
- B. Headache attacks last 4–72 hours (untreated or unsuccessfully treated)
- C. Headache has at least 2 of the following characteristics:
  - 1. Unilateral location
  - 2. Pulsating quality
  - 3. Moderate or severe pain intensity
  - 4. Aggravation by, or causing avoidance of, routine physical activity (e.g., walking or climbing stairs)
- D. During the headache attack, at least 1 of the following:
  - 1. Nausea and/or vomiting
  - 2. Photophobia and phonophobia
- E. Symptoms not attributed to another disorder

**ICHD-2 diagnostic criteria for 1.2 migraine with aura (modified)**

- A. At least 2 attacks fulfilling criteria B–D
- B. Aura consists of at least 1 of the following:
  - 1. Fully reversible visual symptoms
  - 2. Fully reversible sensory symptoms
  - 3. Fully reversible dysphasic speech disturbance (not motor weakness)
- C. At least 2 of the following:
  - 1. Homonymous visual symptoms and/or unilateral sensory symptoms
  - 2. At least one aura symptom develops gradually over more than 5 minutes and/or different aura symptoms occur in succession over 25 minutes
- D. Headache fulfills criteria B–D for 1.1 migraine without aura
- E. Symptoms not attributed to another disorder

sometimes occur with nonmigraine headaches and have been reported with cluster headache, chronic paroxysmal hemicranias, and hemicrania continua.

Cerebrovascular disease, especially transient ischemic attacks, may closely resemble the aura of migraine. Features that differentiate the two entities are the aura's slow evolution and the positive nature of the visual and sensory symptoms (e.g., scintillating scotomas rather than blindness). Investigations to rule out cerebrovascular disease are warranted if the symptoms are of rapid onset (less than 5 minutes), the visual or sensory symptoms are negative (blindness or sensory loss), the aura is prolonged (more than 1 hour), or the patient has risk factors for vascular disease including onset after the age of 50 [5].

### Subtypes of migraine

There are several unusual forms of migraine named for the area of the affected nervous system. In all cases, the manifestation of these unusual auras is reversible.

*Familial hemiplegic migraine* (FHM) (ICHD-2 1.2.4), as the name implies, is manifested by hemiparesis sometimes with hemisensory impairment. Ataxia or other features of basilar-type migraine may occur. Unlike a typical aura, the hemiparesis may last for hours. This is the first headache syndrome linked to genetic polymorphisms. The loci for the condition are on chromosomes 1, 2, or 19 [6]. One or more first-degree relatives have similar attacks. People without a family history but otherwise meeting the criteria are classified as having sporadic hemiplegic migraine. The genetics of FHM are discussed in detail in Chapter 6.

In patients with *basilar-type migraine* (ICHD-2 1.2.6), the symptoms associated with the headache implicate the posterior fossa. The ICHD-2 classification requires at least two of the following aura symptoms: vertigo, tinnitus, decreased hearing, double vision, visual defects in both the temporal and nasal fields of both eyes, ataxia, dysarthria, bilateral paresthesias, or decreased consciousness.



*Retinal migraine* (ICHD-2 1.4) is manifested by scintillation, other positive visual symptoms, or more likely scotomas or partial or complete blindness limited to one eye (in contrast to common homonymous symptoms). Many of these patients experience a permanent partial defect representing retinal infarction [7]. Other causes of monocular visual loss (transient ischemic attacks, optic neuropathy, and retinal detachment) must be considered.

*Childhood periodic syndromes* are often precursors of migraine. *Cyclic vomiting* (ICHD-2 1.3.1) is manifested by intractable recurrent vomiting lasting for hours or days. Pallor and lethargy are usually associated. There is no evidence of gastrointestinal disease. Patients with *abdominal migraine* (ICHD-2 1.3.2) have attacks of abdominal pain associated with anorexia or nausea, with or without vomiting, and pallor. The ICHD-2 criteria require at least two of the three associated symptoms. Again, examination and studies rule out gastrointestinal disease. In patients with *benign paroxysmal vertigo of childhood* (ICHD-2 1.3.3), there are recurrent attacks (at least five) of severe vertigo that may last for minutes to hours; the attacks resolve spontaneously. Neurologic and vestibular functions are normal between attacks, as are imaging studies and electroencephalography. When this condition occurs after childhood, the patients often have bouts of migraine during or independent of vertigo. “Migrainous vertigo” or “vertiginous migraine” are not accepted entities in the ICHD-2 classification. Benign recurrent vertigo may occur at any age and often coexists with migraine. It is considered a comorbid condition.

### Complications of migraine

*Chronic migraine* (ICHD-2 1.5.1) is defined as migraine headaches (usually without aura) or TTHs recurring on more than 15 days per month for more than 3 months [8]. At least 50% of the headaches should be migraine based on the presence of migraine features or response to migraine treatment. Chronic migraine evolves from episodic migraine. Many risk factors contribute to the evolution of chronic migraine from episodic migraine (see Chapter 5). Risk factors include frequent episodic headaches, high levels of disability, allodynia, depression, traumatic head injury, and overuse of drugs, most importantly opioid- and barbiturate-containing analgesics. The diag-

nosis of chronic migraine in the setting of medication overuse is complex (see Chapter 10). In practice, we treat medication overuse as a modifier, i.e., chronic migraine with medication overuse.

*Status migrainosus* (ICHD-2 1.5.2) is a term used for a severe and disabling migraine that lasts for more than 3 days. *Prolonged aura* is an extension of the patient’s typical aura for more than 1 hour but less than 1 week. *Persistent aura without infarction* (ICHD-2 1.5.3) refers to persistence of an aura for more than 1 week. The aura is otherwise typical of past attacks. Evaluations do not reveal cerebral infarction. *Migrainous infarction* (ICHD-2 1.5.4) rarely occurs. The typical aura symptom or symptoms persist beyond 1 hour, and neurologic examination or neuroimaging or both confirms cerebral infarction. Other causes of stroke must be excluded.

Many patients exhibit all but one of the criteria for migraine. The term “probable migraine” (ICHD-2 1.6) is then appropriate.

### TTH (ICHD-2 2.0)

This term replaces the former names of “tension headache” (implying emotional stress) and “muscle contraction headache” (implying a muscular origin of the headache). Although TTH is the most common primary headache, its mechanism is the least well understood. Many believe that TTHs are fragments of migraine. The diagnostic criteria are listed in Table 1.3. The headache characteristics are purposely nonmigrainous. The location is bilateral, the quality is nonpulsating, the intensity is mild to moderate, and the headache is not aggravated by physical activity. Similarly, there is a paucity of associated symptoms: no nausea or vomiting, although either photophobia or phonophobia may be present, but not both. There may or may not be associated pericranial tenderness.

TTHs are classified into episodic (fewer than 15 attacks per month) and chronic (more than 15 attacks per month). Episodic TTH is further subdivided into *infrequent episodic TTH* (ICHD-2 2.1) (less than 1 day per month) and *frequent episodic TTH* (ICHD-2 2.2) (1–14 days per month). *Chronic TTH* (ICHD-2 2.3) usually evolves from the episodic form and in its purest form should not be diagnosed in patients overusing acute medication. In practice, patients with medication overuse headache (a secondary headache)

**Table 1.3.** ICHD-2 Diagnostic criteria for 2.0 tension-type headache (TTH) (modified)

- A. Episodes occurring in frequencies below (see 2.1, 2.2 and 2.3A)
  - B. Headache lasting from 30 minutes to 7 days
  - C. Headache has at least 2 of the following characteristics:
    1. Bilateral location
    2. Pressing/tightening (non-pulsating) quality
    3. Mild or moderate intensity
    4. Not aggravated by routine physical activity, such as walking or climbing upstairs
  - D. Both of the following:
    1. No nausea or vomiting (anorexia may occur)
    2. No more than one of the following: photophobia or phonophobia
  - E. Symptoms not attributed to another disorder
- 2.1 Infrequent episodic TTH  
At least 10 episodes occurring on <1 day per month on average (<12 days per year) and fulfilling criteria B–D
- 2.2 Frequent episodic TTH  
At least 10 episodes occurring on  $\geq 1$  but <15 days per month for at least 3 months ( $\geq 12$  and <180 days per year) and fulfilling criteria B–D
- 2.3 Chronic TTH
- A. Headache occurring on >15 days per month on average for >3 months (>180 days per year) and fulfilling criteria C–E (above)
  - B. Headache lasts hours or may be continuous

usually manifest the features of chronic TTH. As with the classification of migraine, the diagnosis of *probable TTH* (ICHD-2 2.4) is warranted if all but one of the ICHD-2 criteria are met. Another headache that phenotypically may resemble chronic TTH is new daily persistent headache (see “Other primary headaches”).

### TACs (ICHD-2 2.3.0)

The group of primary headaches known as TACs are characterized by trigeminal activation with unilateral pain typically affecting one orbit and ipsilateral autonomic activation (predominantly parasympathetic) [9]. The pain may spread to adjacent areas of one side and is usually locked to the same side on repeated attacks.

*Cluster headache* (ICHD-2 3.1) is the best known of the TACs, and the criteria for diagnosing it are noted in Table 1.4. The pain, usually in or around one orbit, is excruciating and is typically boring, sharp, or stabbing. It begins quickly and lasts from 15 minutes to 3 hours. During the attack, one or more of the following associated features are present: ipsilateral redness and tearing of the eye, clogging or drainage from the nostril, and ptosis and miosis. The attacks may occur between one and eight times per day, often

**Table 1.4.** ICHD-2 diagnostic criteria for cluster headache

- 3.0 Cluster headache (modified)
- A. At least 5 attacks fulfilling criteria B–D
  - B. Severe or very severe unilateral orbital, supraorbital and/or temporal pain lasting 15–180 minutes if untreated
  - C. Headache is accompanied by at least 1 of the following:
    1. Ipsilateral conjunctival injection and/or lacrimation
    2. Ipsilateral nasal congestion and/or rhinorrhea
    3. Ipsilateral eyelid edema
    4. Ipsilateral forehead and facial sweating
    5. Ipsilateral miosis and/or ptosis
    6. A sense of restlessness or agitation
  - D. Attacks have a frequency from one every other day to 8 per day
  - E. Not attributed to another disorder
- 3.1 Episodic cluster headache occurs in periods lasting 7 days to 1 year
- 3.2 Chronic cluster headache occurs for more than 1 year without remission

awakening the patient from sleep. During the headache, in contrast to the behavior during migraine, the patient cannot lie still but is impelled to move about or sit and rock. Cluster headaches affect men more than women. These people are often heavy smokers.

The most common subtype is *episodic cluster headache* (ICHD-2 3.1.1), recurring every day for weeks or months (in clusters), and then remitting for one or more months annually, often recurring during the same season or seasons every year. During these cluster periods, alcoholic beverages and other vasodilating agents may trigger an attack. The unfortunate individuals who do not experience a remission within a year or have only a brief remission of less than 1 month are classified as having *chronic cluster headache* (ICHD-2 3.1.2).

*Paroxysmal hemicranias*, as the name implies, are unilateral headaches that are of short duration (2–30 minutes) and occur more than five times per day [9]. Associated with these are some or all of the autonomic features of cluster headache. The headaches invariably respond to prophylactic doses of indomethacin. The headaches usually occur without remission: *chronic paroxysmal hemicrania* (ICHD-2 3.2.2). Less frequent are remissions of 1 month or more: *episodic paroxysmal hemicrania* (ICHD-2 3.2.1).

*Short-lived unilateral neuralgiform headache attacks with conjunctival injection and tearing (SUNCT) syndrome* (ICHD-2 3.3)—the name of this syndrome tells it all [10]. The headache and autonomic features are again similar to those of cluster headaches and paroxysmal hemicranias, but the duration is shorter (5 seconds to 4 minutes) and the frequency is higher (3–200 attacks per day).

### Other primary headaches

*Hemicrania continua* (ICHD-2 4.7) is a daily and continuous unilateral headache that always responds to therapeutic doses of indomethacin [11]. The daily pain is of moderate intensity, but superimposed exacerbations often occur and may mimic migraine or cluster headache. Patients sometimes complain only of the migraine or cluster headache symptoms. A sensation of conjunctival irritation is often noted. Without inquiring about lesser daily headaches, the diagnosis of hemicrania continua may be missed.

### ★ TIPS AND TRICKS

Patients often describe the headache that is most intense or disabling, such as migraine. Ask whether or not some headache or discomfort is present between the severe attacks. This may lead to the diagnosis of hemicrania continua with exacerbations that resemble a migraine-like headache.

*Primary stabbing headaches* (ICHD-2 4.1) are manifested by episodes of localized stabs of pain [12]. Formerly called “jabs and jolts,” the pains last only one or a few seconds and occur at random without a consistent location. The frequency may range from a few per day to a few per month. Associated symptoms are lacking. These headaches are not uncommon in people with migraine.

Other primary headaches may be triggered by *cough, exertion, or sexual activity* (ICHD-2 4.2, ICHD-2 4.3, ICHD-2 4.4) and are labeled as such. Organic disease may cause headache evoked by these activities. Neuroimaging with special attention to the posterior fossa is mandatory to rule out cerebral aneurysm, subarachnoid hemorrhage, arterial dissection, and Arnold–Chiari malformation. Headache associated with sexual activity may be preorgasmic, causing a dull ache in the head and neck, or orgasmic, which often has an explosive quality.

*Primary thunderclap headache* (ICHD-2 4.6) is a severe headache of sudden onset [13]. The attack mimics the pain of subarachnoid hemorrhage, which must be excluded. The headache lasts from 1 to 10 days and may recur at random over weeks or months.

*Hypnic headaches* (ICHD-2 4.7) awaken patients from sleep, often at the same time each night. The pain is not specific and is without associated symptoms. Usually mild to moderate in intensity, the headache lasts about 30 minutes. It occurs mainly in late middle age or beyond.

*New daily persistent headache* (ICHD-2 4.8) was initially thought to mainly resemble chronic TTH. Recent evaluation, however, has revealed that most people with this condition have many features of migraine, and a modified classification has been proposed [14]. Patients can often

identify the exact day or days of onset. The headaches begin within 3 days of the noted day and are constant for more than 3 months. Acute medication overuse must be first considered, and chronic migraine should also be ruled out before the diagnosis is made.

### Headache diagnosis

In beginning to formulate a diagnosis, the few steps in the algorithm of Figure 1.1 should be helpful. As noted before, headaches secondary to organic disease or dysfunction must be ruled out

before a diagnosis of primary headache disorder is established. Neurologic workup is not necessary with every new patient, but if the headache is atypical or has recently changed or there is a history of red flags, a search for underlying disease is warranted (Table 1.5).

In formulating the differential diagnoses of the primary headaches, it is useful to follow a sequence of steps (Table 1.6). First, classify the headaches into low or high frequency (less or more than 15 headache days per month). Second, divide the headaches into those of short or long

**Table 1.5.** Alarms warranting further consideration

| Red flag  | Consider  | Possible investigation(s)  |
|---|---|--|
| Sudden-onset headache   | Subarachnoid hemorrhage, bleed into mass or AVM, mass lesion (especially posterior fossa), pituitary apoplexy | Neuroimaging<br>Lumbar puncture (after neuroimaging evaluation)                |
| Worsening-pattern of headache (change in frequency or character)              | Subdural hematoma, other mass lesion, medication overuse  | Neuroimaging   |
| Headache with systemic illness (fever, neck stiffness, rash)                  | Meningitis, encephalitis, Lyme disease, systemic infection, collagen vascular disease, arteritis              | Neuroimaging<br>Lumbar puncture<br>Biopsy<br>Blood tests                       |
| Focal neurologic signs, or symptoms other than typical visual or sensory aura | Mass lesion, AVM, collagen vascular disease   | Neuroimaging<br>Lumbar puncture (after neuroimaging evaluation)<br>Blood tests |
| Papilledema   | Mass lesion, idiopathic intracranial hypertension, encephalitis, meningitis                                   | Neuroimaging<br>Lumbar puncture (after neuroimaging evaluation)                |
| Triggered by cough, exertion or sexual activity                               | Subarachnoid hemorrhage, mass lesion  | Neuroimaging<br>Consider lumbar puncture                                       |
| Headache during pregnancy or post-partum                                      | Cortical vein/cranial sinus thrombosis, carotid dissection, pituitary apoplexy                                | Neuroimaging   |
| New headache type in a patient with cancer                                    | Metastasis  | Neuroimaging<br>Lumbar puncture  |
| Lyme disease  | Meningoencephalitis   | Neuroimaging<br>Lumbar puncture  |
| HIV   | Opportunistic infection, tumor  | Neuroimaging<br>Lumbar puncture  |
| Age of onset over 50  | Organic disease: e.g., giant cell arteritis, mass lesion  | Neuroimaging<br>Erythrocyte sedimentation rate                                 |

AVM, arteriovenous malformation.

**Table 1.6.** Primary headache diagnosis by syndromic group

| Frequency                               | Short duration (<4 hours)  | Long duration (4 hours or more)   |
|---|--|---|
| Low to moderate<br>(<15 days per month) | Primary stabbing headache<br>Thunderclap headache<br>SUNCT syndrome<br>Hypnic headache   | Migraine<br>Episodic TTH  |
| Chronic (15 or more<br>days per month)  | Cluster headache and other TACs<br>Paroxysmal hemicrania<br>Primary stabbing headache<br>Thunderclap headache<br>SUNCT syndrome<br>Hypnic headache | Chronic migraine<br>Chronic TTH<br>Hemicrania continua<br>New daily persistent headache |

duration (less than or more than 4 hours). Finally, consider headaches of short duration of low or high frequency and the presence or not of triggering factors. This approach should lead to the correct diagnosis for almost all patients.

## References

- Ad Hoc Committee on Classification of Headaches. Classification of headache. *JAMA* 1962;**179**:717–8.
- Headache Classification Committee of the International Headache Society. Classification and diagnostic criteria for headache disorders, cranial neuralgias and facial pain. *Cephalalgia* 1988;**8**(Suppl. 7):1–96.
- Headache Classification Committee of the International Headache Society. The International Classification of Headache Disorders. *Cephalalgia* 2004;**24**:1–160.
- Lipton RB, Stewart WF, Diamond S, et al. Prevalence and burden of migraine in the United States: data from the American Migraine Study II. *Headache* 2001;**41**:646–57.
- Fisher CM. Late life migraine accompaniments as a cause of unexplained transient ischemic attacks. *Can J Neurol Sci* 1980;**7**:9–17.
- Ducros A, Denier C, Joutel A, et al. The clinical spectrum of familial hemiplegic migraine associated with mutations in a neuronal calcium channel. *N Engl J Med* 2001;**345**:17–24.
- Grosberg BM, Solomon S, Friedman DI, et al. Retinal migraine reappraised. *Cephalalgia* 2006;**26**:1275–86.
- Silberstein SD, Lipton RB, Sliwinski M. Classification of daily and near daily headaches: field trial of revised IHS criteria. *Neurology* 1996;**47**:871–5.
- Goadsby PJ, Lipton RB. A review of paroxysmal hemicranias, SUNCT syndrome and other short-lasting headaches with autonomic features, including new cases. *Brain* 1997;**120**:193–209.
- Sjaastad O, Saunte C, Salvesen R, et al. Short-lasting unilateral neuralgiform headache attacks with conjunctival injection, tearing, sweating, and rhinorrhea. *Cephalalgia* 1989;**9**:147–56.
- Sjaastad O, Spierings EL. Hemicrania continua: another headache absolutely responsive to indomethacin. *Cephalalgia* 1984;**4**:65–70.
- Pareja JA, Rujiz J, Deisla C, et al. Idiopathic stabbing headache (jabs and jolt syndrome). *Cephalalgia* 1996;**16**:93–6.
- Day JW, Raskin NH. Thunderclap headache: symptom of unruptured cerebral aneurysm. *Lancet* 1986;**2**:1247–8.
- Robbins MS, Grosberg BM, Napchan U, et al. Clinical and prognostic subforms of new daily persistent headache. *Neurology* 2010;**74**:1358–64.
- Lipton RB, Silberstein SD, Dodick D. Overview of diagnosis and classification. In Silberstein SD, Lipton RB, & Dodick DW (eds.), *Wolff's Headache and Other Head Pain* (8th edn). New York: Oxford University Press; 2008: p. 38.

# Approach to the Patient with Headache

**Brian M. Grosberg, Benjamin W. Friedman, and Seymour Solomon**

Albert Einstein College of Medicine, Bronx, NY, USA

## Introduction

Headache is one of the most common complaints presenting to primary care physicians and neurologists. Although the vast majority of headache syndromes are benign, clinicians are faced with the crucial task of differentiating benign headache disorders from potentially life-threatening conditions. Given the broad range of disorders that present with headache, a focused and systematic approach is necessary to facilitate the prompt diagnosis and treatment of various kinds of head pain.

In managing a patient with chronic, recurrent primary headaches, there must be a positive interrelationship between patient and doctor. Some patients sit back and say “I’m here. It’s your job to cure me.” Some doctors quickly prescribe an analgesic and give the patient short shrift. During the initial evaluation, the patient and the doctor should get to know one another. Patients want their doctor to take an interest in them as a person. They want to know why they have a headache and what is causing it. The doctor wants not only to make a diagnosis, but also to judge the patient’s personality and needs. Effective communication and interpersonal skills are integral to the care of headache patients [1].

There are several steps in patient management. First, of course, is the history and examination. One must differentiate primary from secondary headaches. If worrisome features or “red flags” are present either in the history or on examination, diagnostic testing may be necessary to exclude secondary headache disorders. A summary of these red flags is listed in Table 2.1 [2]. Having determined that the condition is a primary headache, the specific headache type must be diagnosed. Once a diagnosis has been established, the general therapeutic options should be outlined in a treatment plan tailored to the patient. By taking into account the patient’s headache diagnosis or diagnoses, frequency of headache attacks, headache-related functional impairment, comorbid medical conditions, frequency of acute medication use, health-related quality of life, and treatment preferences, clinicians can then develop an optimal treatment plan for each patient. One should emphasize the need for the patient to be active in his or her own care. Realistic expectations should be established, and rarely will the initial plan be perfect. Follow-up visits are scheduled to modify the therapy and, if necessary, to reassess the diagnosis.

*Headache*, First Edition. Edited by Matthew S. Robbins, Brian M. Grosberg, and Richard B. Lipton.  
© 2013 John Wiley & Sons, Ltd. Published 2013 by John Wiley & Sons, Ltd.