

Asymmetric Phase Transfer Catalysis

Edited by
Keiji Maruoka



WILEY-
VCH

WILEY-VCH Verlag GmbH & Co. KGaA

Asymmetric Phase Transfer Catalysis

Edited by

Keiji Maruoka

Related Titles

M. Christmann, S. Bräse (Eds.)

Asymmetric Synthesis – The Essentials

2006

ISBN: 978-3-527-31399-0

B. Cornils, W.A. Herrmann, I.T. Horvath, W. Leitner,
S. Mecking, H. Olivier-Bourbigou, D. Vogt (Eds.)

Multiphase Homogeneous Catalysis

2005

ISBN: 978-3-527-30721-0

J. Christoffers, A. Baro (Eds.)

Quaternary Stereocenters

Challenges and Solutions for Organic Synthesis

2005

ISBN: 978-3-527-31107-1

A. Berkessel, H. Gröger

Asymmetric Organocatalysis

**From Biomimetic Concepts to Applications
in Asymmetric Synthesis**

2005

ISBN: 978-3-527-30517-9

Asymmetric Phase Transfer Catalysis

Edited by
Keiji Maruoka



WILEY-
VCH

WILEY-VCH Verlag GmbH & Co. KGaA

The Editor

Prof. Keiji Maruoka

Kyoto University
Department of Chemistry
Graduate School of Science
Sakyo
Kyoto 606-8502
Japan

All books published by Wiley-VCH are carefully produced. Nevertheless, authors, editors, and publisher do not warrant the information contained in these books, including this book, to be free of errors. Readers are advised to keep in mind that statements, data, illustrations, procedural details or other items may inadvertently be inaccurate.

Library of Congress Card No.: applied for

British Library Cataloguing-in-Publication Data

A catalogue record for this book is available from the British Library

**Bibliographic information published by
the Deutsche Nationalbibliothek**

Die Deutsche Nationalbibliothek lists this publication in the Deutsche Nationalbibliografie; detailed bibliographic data are available in the Internet at <http://dnb.d-nb.de>

© 2008 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

All rights reserved (including those of translation into other languages). No part of this book may be reproduced in any form – by photoprinting, microfilm, or any other means – nor transmitted or translated into a machine language without written permission from the publishers. Registered names, trademarks, etc. used in this book, even when not specifically marked as such, are not to be considered unprotected by law.

Printed in the Federal Republic of Germany
Printed on acid-free paper

Cover design Adam Design, Weinheim

Typesetting Thomson Digital, Noida, India

Printing Strauss GmbH, Mörlenbach

Bookbinding Litges & Dopf GmbH, Heppenheim

ISBN: 978-3-527-31842-1

Contents

Preface XI

List of Contributors XIII

1	The Basic Principle of Phase-Transfer Catalysis and Some Mechanistic Aspects	1
	<i>Takuya Hashimoto and Keiji Maruoka</i>	
1.1	Introduction	1
1.2	Inorganic Base-Promoted Activation of Acidic Organic Compounds	2
1.2.1	Generation of Reactive Onium Carbanion Species	2
1.2.2	Stability of the Onium Carbanion	3
1.2.3	Reactivity of the Onium Carbanion	3
1.2.4	Nucleophilic Substitution Reaction	4
1.2.5	Nucleophilic Addition to Electrophilic C=X Double Bonds	5
1.3	Phase-Transfer-Catalyzed Addition of Anion Supplied as Metal Salt	6
1.4	Use of Crown Ether as Phase-Transfer Catalyst	7
	References	7
2	Cinchona-Derived Chiral Phase-Transfer Catalysts for Amino Acid Synthesis	9
	<i>Takashi Ooi</i>	
2.1	Introduction	9
2.2	α -Amino Acid Synthesis	9
2.2.1	Monoalkylation of Schiff Bases Derived from Glycine	9
2.2.2	Alkylation of Schiff Bases Derived from α -Alkyl- α -Amino Acids	19
2.2.3	Other Alkylations for α -Amino Acid Synthesis	21
2.2.4	Michael Reaction of Glycinate Benzophenone Schiff Bases	22

2.2.5	Aldol and Related Reactions	24
2.2.6	Aza-Henry Reaction	25
2.2.7	Strecker Reaction	27
2.2.8	Aziridination	27
2.2.9	Radical Reaction	28
2.3	β -Amino Acid Synthesis	29
2.3.1	Mannich Reaction	29
2.3.2	Cyclopropanation	29
2.4	Conclusions	30
	References	31

3 Cinchona-Derived Chiral Phase-Transfer Catalysts for Other Asymmetric Synthesis 35

Shigeru Arai

3.1	Introduction	35
3.2	Asymmetric Darzens Reaction	35
3.3	Asymmetric Conjugated Addition	36
3.4	Asymmetric Aldol Reaction	39
3.5	Asymmetric Oxygen-Functionalization	40
3.6	Asymmetric Alkylation	42
3.7	Asymmetric Alkenylation and Alkynylation	43
3.8	Asymmetric S_N Aromatic Reaction	45
3.9	Asymmetric Strecker Synthesis	46
3.10	Asymmetric Fluorination	46
	References	47

4 Cinchona-Derived Chiral Poly(Phase-Transfer Catalysts) for Asymmetric Synthesis 49

Sang-sup Jew and Hyeung-geun Park

4.1	Cinchona Alkaloids	49
4.1.1	Cinchona Alkaloids in Asymmetric Phase-Transfer Catalysis	49
4.1.2	The Origin of Stereoselectivity of Cinchona-PTCs	50
4.2	Development of Dimeric Cinchona-PTCs by the Park-Jew Group	51
4.2.1	Dimeric Cinchona-PTCs with Phenyl Linker	51
4.2.2	1,3-Dimeric Cinchona-PTCs with Electronically Modified Phenyl Linker	56
4.2.3	Polymeric Cinchona-PTCs with Other Linkers	56
4.2.4	1,3-Phenyl- and 2,7-Naphthyl-Linked Dimeric Cinchona-PTCs	57
4.3	Polymeric PTCs Developed by the Najera Group	61
4.4	Polymeric PTCs Developed by the Siva Group	62
4.5	Polymeric PTCs Developed by the Wang Group	63
4.6	Asymmetric Epoxidation with Polymeric Cinchona-PTCs	63
4.7	Conclusions	68
	References	68

5	Binaphthyl- and Biphenyl-Modified Chiral Phase-Transfer Catalysts for Asymmetric Synthesis	71
	<i>Keiji Maruoka</i>	
5.1	Introduction	71
5.2	Alkylation	71
5.2.1	Asymmetric Synthesis of α -Alkyl α -Amino Acids and Their Derivatives	71
5.2.1.1	Asymmetric Monoalkylation of Glycine Ester Schiff Bases	71
5.2.1.2	Asymmetric Monoalkylation of Glycine Amide Schiff Bases	85
5.2.1.3	Diastereoselective Alkylation of Glycine Schiff Base with Optically Enriched Alkyl Halides	87
5.2.1.4	Recyclable Catalysts and Reagents and Solid-Phase Synthesis	87
5.2.1.5	Application of Asymmetric Synthesis of α -Amino Acids	87
5.2.2	Asymmetric Synthesis of α,α -Dialkyl- α -Amino Acids	90
5.2.3	Alkylation of Schiff Base-Activated Peptides	94
5.2.4	Other Alkylations	97
5.3	Michael Addition	99
5.4	Aldol Reaction	105
5.5	Mannich Reaction	106
5.6	Neber Rearrangement	106
5.7	Epoxidation	108
5.8	Strecker Reaction	109
5.9	Conclusions	110
	References	111
6	Two-Center Chiral Phase-Transfer Catalysts for Asymmetric Synthesis	115
	<i>Masakatsu Shibasaki and Takashi Ohshima</i>	
6.1	Introduction	115
6.2	Design and Synthesis of Two-Center Chiral Phase-Transfer Catalyst	115
6.3	Catalytic Asymmetric Phase-Transfer Alkylation	118
6.4	Catalytic Asymmetric Phase-Transfer Michael Addition to α,β -Unsaturated Esters	119
6.5	Catalytic Asymmetric Phase-Transfer Michael Addition to Enones	120
6.6	Catalytic Asymmetric Phase-Transfer Mannich-Type Reaction	122
6.7	Synthetic Applications	123
6.7.1	Enantioselective Syntheses of Aeruginosin 298-A and its Analogues	123
6.7.2	Short Syntheses of (+)-Cylindricine and Formal Synthesis of (–)-Lepadiformine	124
6.7.3	Formal Synthesis of (+)-CP-99,994 and Total Synthesis of (+)-Nemonapride	125
6.8	Catalyst Recovery and Reuse	126
6.9	Role of Two Ammonium Cations	127
6.10	Other Two-Center Chiral Phase-Transfer Catalysts	128
6.10.1	Two-Center Chiral Phase-Transfer Catalyst Derived from BINOL (1)	128

6.10.2	Two-Center Chiral Phase-Transfer Catalyst Derived from BINOL (2)	129
6.10.3	Two-Center Chiral Phase-Transfer Catalyst Derived from BINOL (3)	129
6.10.4	Two-Center Chiral Phase-Transfer Catalyst Derived from Tartrate (1)	130
6.10.5	Two-Center Chiral Phase-Transfer Catalyst Derived from Tartrate (2)	132
6.11	Conclusions	132
	References	133

7 Other Chiral Phase-Transfer Catalysts for Asymmetric Synthesis 135

Hiroaki Sasai and Mahesh L. Patil

7.1	Introduction	135
7.2	Other Chiral Phase-Transfer-Catalyzed Reactions	136
7.2.1	Alkylation	136
7.2.1.1	Substrate Scope	147
7.2.1.2	Mechanistic Investigations	149
7.2.2	Michael Reaction	150
7.2.3	Epoxidation	152
7.2.4	Hydroxylation	156
7.3	Conclusions	157
	References	157

8 Crown Ethers, Taddol, Nobin and Metal(salen) Complexes as Chiral Phase-Transfer Catalysts for Asymmetric Synthesis 161

Michael North

8.1	Introduction	161
8.2	Crown Ethers as Chiral Phase-Transfer Catalysts	163
8.2.1	Use of Crown Ethers in Asymmetric Michael Additions	163
8.2.2	Use of Crown Ethers in Darzens Condensations	165
8.2.3	Use of Crown Ethers in Asymmetric Epoxidations	166
8.3	Use of Taddolates as Chiral Phase-Transfer Catalysts	166
8.4	Use of Nobin and Related Species as Asymmetric Phase-Transfer Catalysts	170
8.5	Use of Metal(salen) Complexes as Chiral Phase-Transfer Catalysts	173
8.5.1	Nickel(salen) Complexes	173
8.5.2	Copper(salen) Complexes	175
8.5.3	Use of Other Metal(salen) Complexes as Catalysts for Enolate Alkylation	182
8.5.4	Metal(salen) Complexes as Catalysts for Darzens Condensations	183
8.6	Conclusions	184
	References	185

9	Chiral Quaternary Ammonium Fluorides for Asymmetric Synthesis	189
	<i>Seiji Shirakawa, Takashi Ooi, and Keiji Maruoka</i>	
9.1	Introduction	189
9.2	<i>In-Situ</i> Generation of Chiral Quaternary Ammonium Fluorides	189
9.3	Chiral Quaternary Ammonium Fluorides: Preparation and Application to Organocatalytic Asymmetric Reactions	193
9.4	Chiral Quaternary Ammonium Bifluorides: Preparation and Use as Organocatalysts for Asymmetric Carbon–Carbon Bond-Forming Reactions	197
9.5	Conclusions	205
	References	205
	Index	207

Preface

In contrast to the wide availability of phase-transfer reactions and processes using non-chiral phase-transfer catalysts, the development of asymmetric phase-transfer catalysis based on the use of structurally well-defined chiral, non-racemic catalysts has progressed rather slowly, despite the great importance of creating a new domain in modern asymmetric catalysis by taking full advantage of structurally and stereochemically modifiable tetra-alkylammonium salts. Following the first meaningful discovery of *N*-(*p*-(trifluoromethyl)benzyl)cinchonidium bromide as chiral phase-transfer catalyst by the Merck group in 1984, various types of cinchona-alkaloid-derived chiral phase-transfer catalysts have been developed. The major applications of such catalysts include the enantioselective synthesis of amino acid derivatives by the asymmetric alkylation of glycine and other α -alkylamino acid derivatives. Although O'Donnell, in 1989, applied *N*-(benzyl)cinchoninium chloride and its cinchonidine derivative to the asymmetric synthesis of both enantiomers of α -alkylamino acid derivatives, research in this area subsequently made little progress for some time after these milestone reports. However, in 1997 a new class of cinchona alkaloid-derived catalysts bearing an *N*-anthracenylmethyl function, and developed independently by Corey and Lygo, opened a new era of asymmetric phase-transfer catalysis.

Two years later, in 1999, a totally new aspect on the design of chiral phase-transfer catalysts was reported, based on the use of non-cinchona-alkaloid derivatives. These structurally rigid, chiral spiro ammonium salts were introduced by using commercially available (*S*)- or (*R*)-1,1'-bi-2-naphthol as a new C_2 -symmetric chiral phase-transfer catalyst, and successfully applied to the highly efficient, catalytic enantioselective alkylation of glycine Schiff base under mild phase-transfer conditions. The rational design of chiral binaphthyl-modified phase-transfer catalysts was relatively straightforward in comparison with cinchona-alkaloid-derived chiral phase-transfer catalysts, and was realized by fine-tuning the 3,3'-diaryl substituents of the chiral binaphthyl moiety. This allowed the systematic development of a variety of new – yet practical – asymmetric transformations under mild phase-transfer conditions, and inspired the further design of cinchona alkaloid-derived phase-transfer catalysts and other forms of chiral phase-transfer catalyst. Accordingly,

major efforts towards this direction have recently resulted in notable achievements, whereby it has become feasible to perform a variety of enantioselective bond-formation reactions under mild phase-transfer-catalyzed conditions.

The aim of this book is to provide a concise and comprehensive treatment of this continuously growing field of catalysis, focusing not only on the design of the various types of chiral phase-transfer catalyst but also on the synthetic aspects of this chemistry. In addition, the aim is to promote the synthetic applications of these asymmetric phase-transfer reactions by giving solid synthetic evidence. Clearly, despite recent spectacular advances in this area, there is still plenty of room for further continuous development in asymmetric phase-transfer catalysis.

Kyoto, January 2008

Keiji Maruoka

List of Contributors

Shigeru Arai

Chiba University
Graduate School of Pharmaceutical
Sciences
1-33 Yayoi-cho, Inage-ku
Chiba, 263-8522
Japan

Takuya Hashimoto

Kyoto University
Department of Chemistry
Graduate School of Science
Sakyo
Kyoto 606-8502
Japan

Sang-sup Jew

College of Pharmacy
Seoul National University
San 56-1
Sinlim-dong, Kwanak-gu
Seoul 151-742
South Korea

Keiji Maruoka

Kyoto University
Department of Chemistry
Graduate School of Science
Sakyo
Kyoto 606-8502
Japan

Michael North

University of Newcastle upon Tyne
School of Natural Sciences
Bedson Building
Newcastle upon Tyne, NE1 7RU
United Kingdom

Takashi Ohshima

Osaka University
Department of Chemistry
Graduate School of Engineering Science
Toyonaka
Osaka 560-8531
Japan

Takashi Ooi

Nagoya University
Department of Applied Chemistry
Graduate School of Engineering
Chikusa
Nagoya 464-8603
Japan

Hyeung-geun Park

Seoul National University
College of Pharmacy
San 56-1
Sinlim-dong Kwanak-gu
Seoul 151-742
South Korea

Mahesh L. Patil

Osaka University
The Institute of Scientific and
Industrial Research (ISIR)
8-1 Mihogaoka, Ibaraki-shi
Osaka 567-0047
Japan

Hiroaki Sasai

Osaka University
The Institute of Scientific and
Industrial Research (ISIR)
8-1 Mihogaoka, Ibaraki-shi
Osaka 567-0047
Japan

Masakatsu Shibasaki

The University of Tokyo
Graduate School of Pharmaceutical
Sciences
7-3-1 Hongo, Bunkyo-ku
Tokyo 113-0033
Japan

Seiji Shirakawa

Nihon University
Department of Applied Molecular
Chemistry
College of Industrial Technology
Narashino
Chiba 275-8575
Japan

1


The Basic Principle of Phase-Transfer Catalysis and Some Mechanistic Aspects

Takuya Hashimoto and Keiji Maruoka

1.1

Introduction

In 1971, Starks introduced the term “phase-transfer catalysis” to explain the critical role of tetraalkylammonium or phosphonium salts (Q^+X^-) in the reactions between two substances located in different immiscible phases [1]. For instance, the displacement reaction of 1-chlorooctane with aqueous sodium cyanide is accelerated many thousand-fold by the addition of hexadecyltributylphosphonium bromide **1** as a phase-transfer catalyst (Scheme 1.1). The key element of this tremendous reactivity enhancement is the generation of quaternary phosphonium cyanide, which renders the cyanide anion organic soluble and sufficiently nucleophilic.



Scheme 1.1

Although this was not the first observation of the catalytic activity of quaternary onium salts, the foundations of phase-transfer catalysis were laid by Starks, together with Makosza and Brändström, during the mid to late 1960s [2]. Ever since that time, the chemical community has witnessed the steady growth of phase-transfer catalysis as a practical methodology for organic synthesis, featuring its simple experimental operations, mild reaction conditions, inexpensive and environmentally benign reagents and solvents, and the possibility to conduct large-scale preparations [3,4]. Nowadays, phase-transfer catalysis appears to be a prime synthetic tool, being appreciated not only in various fields of organic chemistry but also among widespread industrial applications.

Despite the development of phase-transfer catalysis in organic synthesis, the mechanistic aspects of phase-transfer catalysis remain obscure, due mainly to the

difficulty of investigating biphasic systems and the many complex parameters that must be analyzed.

The aim of this chapter is to assist the reader to generate an intuitive understanding of the mechanism of asymmetric phase-transfer catalysis, together with a practical guide for the design of such processes. More detailed studies related to the physical and numerical aspects of phase-transfer catalysis may be consulted elsewhere [3].

1.2

Inorganic Base-Promoted Activation of Acidic Organic Compounds

The representative reaction system applied in asymmetric phase-transfer catalysis is the biphasic system composed of an organic phase containing an acidic methylene or methine compound and an electrophile, and an aqueous or solid phase of inorganic base such as alkaline metal (Na, K, Cs) hydroxide or carbonate. The key reactive intermediate in this type of reaction is the onium carbanion species, mostly onium enolate or nitronate, which reacts with the electrophile in the organic phase to afford the product.


1.2.1

Generation of Reactive Onium Carbanion Species

The exact pathway for generating the reactive onium carbanion species remains the subject of controversy, typically among Starks extraction mechanism (Scheme 1.2) and the Makosza interfacial mechanism (Scheme 1.3).


In the Starks extraction mechanism, the phase-transfer catalyst moves back and forth across the organic and aqueous phases. The onium salt equilibrates with the inorganic base in the aqueous phase, and extracts hydroxide into the organic phase.

Starks Extraction Mechanism



Scheme 1.2

Makosza Interfacial Mechanism



Scheme 1.3

The onium hydroxide then abstracts hydrogen from the acidic organic compound to give the reactive intermediate Q^+R^- .

The advocated pathway of the interfacial mechanism is the first formation of metal carbanion at the interface of organic and aqueous phase in the absence of phase-transfer catalyst, followed by the extraction of the formed metal carbanion species from the interface into the organic phase by the action of phase-transfer catalyst.

Since asymmetric phase-transfer catalysts normally contain highly lipophilic chiral organic frameworks, and are reluctant to enter the aqueous phase, the Makosza interfacial mechanism seems plausible.

Clearly, the area of the interface and the basicity of the inorganic salt affect the amount of available onium carbanion. It should be also noted that an excessively lipophilic phase-transfer catalyst would hardly access the interface, and consequently the use of such a catalyst would result in an insufficient reaction.

1.2.2

Stability of the Onium Carbanion


The onium carbanion formed under phase-transfer conditions is unstable depending on the anion source, and in the absence of an electrophilic reaction partner, degradation of the accumulated onium carbanion in the organic phase may be observed. This is known to proceed via Hoffman elimination, nucleophilic substitution and/or Stevens rearrangement (Scheme 1.4) [4f,6,7]. The direct decomposition of onium salt, as influenced by the strong inorganic base at the interface, may be also operative.

1.2.3


Reactivity of the Onium Carbanion

Cation exchange from the metal cation to the onium carbanion improves the intrinsic reactivity of the latter due to formation of the “naked anion”. At the same time, the


Hoffman Elimination



Nucleophilic Substitution



Stevens Rearrangement



Scheme 1.4

onium carbanion in the organic phase is less hydrated compared to the metal carbanion at the interface, which functions as another factor for an enhanced reactivity of the onium carbanion.

1.2.4


Nuclueophilic Substitution Reaction



The fate of the onium carbanion Q^+R^- incorporated into the organic phase depends on the electrophilic reaction partner. The most studied area in the asymmetric phase-transfer catalysis is that of asymmetric alkylation of active methylene or methine compounds with alkyl halides, in an irreversible manner. The reaction mechanism illustrated above is exemplified by the asymmetric alkylation of glycine Schiff base (Scheme 1.5) [8].

In the first step, glycine Schiff base **2** reacts with the inorganic base at the interface of two phases to give the metal enolate **3**, which remains at the interface due to its highly polar character. The metal enolate **3** then exchanges the cation to provide onium enolate **4**. The sufficiently lipophilic **4** then moves into the organic phase to react with alkyl halide. After the reaction, onium halide is regenerated and enters the next catalytic cycle. The key issue to be considered here is the possibility of product racemization and dialkylation. In this example, the basicity of the inorganic base and acidity of the substrate and product, as well as other reaction conditions, are carefully adjusted to circumvent this problem. It should be also noted that control of the *E/Z* geometry of the enolate is apparently critical to the asymmetric induction, although there is no clear evidence about which isomer is the actual reacting species in this case.

Asymmetric Alkylation of Glycine Schiff Base




Scheme 1.5

An asymmetric S_N2' reaction and S_NAr reaction are included as other examples of this category.

1.2.5


Nucleophilic Addition to Electrophilic $C=X$ Double Bonds



Unlike the nucleophilic substitution reactions which generate stable onium halide after the reaction, nucleophilic additions to electrophilic $C=X$ double bonds ($X=C, N, O$) provide rather basic onium anion species as an initial product. If the anion is sufficiently stable under the reaction conditions, onium anion will then exchange the counter ion for the other metal carbanion at the interface to regenerate the reactive onium carbanion Q^+R^- . In another scenario, the basic onium anion may abstract the acidic hydrogen atom of the other substrate to provide Q^+R^- directly. Such a reaction system ideally requires only a catalytic amount of the base although, in general, a substoichiometric or excess amount of the base is used to lead the reaction to completion. An additional feature of this system is the substantial possibility of a retro-process at the crucial asymmetric induction step, which might be problematic in some cases.

The direct asymmetric aldol reaction under phase-transfer conditions is a representative example of this class of phase-transfer reaction, which is known to proceed with a catalytic amount of base and to include an undesired retro-process (Scheme 1.6) [9]. Here, the onium enolate **4** reacts with aldehyde in the organic

Direct Asymmetric Aldol Reaction of Glycine Schiff Base



Scheme 1.6

phase to give the onium alkoxide **7**, which may abstract the hydrogen from the other glycine Schiff base **2** to provide another reactive intermediate onium enolate **4** and the protonated aldol product **6**. It should be noted that this is not the only expected pathway, and there are complex equilibria in this reaction system which, for reasons of clarity, not fully delineated below. The phase-transfer-catalyzed asymmetric Michael addition and the Mannich reaction are other typical examples which fall into this category.

1.3

Phase-Transfer-Catalyzed Addition of Anion Supplied as Metal Salt

The common feature in the asymmetric phase-transfer catalysis introduced above is formation of the reactive onium carbanion from the inorganic base and active methylene or methine compounds, followed by extraction of the carbanion species as the onium salt from the interfacial area into the organic phase. Thus, both reagents normally remain in the organic phase. In the other category of asymmetric phase-transfer catalysis, the anion itself supplied as an aqueous solution or solid of its inorganic salt as the reaction partner, and the two reactants are strictly separated by the interface. In such a system, the anion is transferred gradually from the aqueous phase into the organic phase by the intervention of an onium salt.


With regards to the mechanism of the generation of onium anion, the Starks extraction mechanism and interfacial mechanism (Brändström–Montanari modification) are suggested (Scheme 1.7). As in the above-described case, the interfacial mechanism seems to be operative in the asymmetric phase-transfer catalysis.

The reaction conditions are rather mild, so that the possibility of side reactions, such as catalyst decomposition, is considerably reduced. The major challenge associated with these reactions is an absence of prochirality in the anionic species. Namely, the chiral onium anion must discriminate the enantiotopic face of the distant electronically neutral reaction partner in the organic phase.


Asymmetric phase-transfer catalyzed oxidation of olefin using sodium hypochlorite or potassium permanganate as metal anion sources is the typical example of this category.

The asymmetric epoxidation of an α,β -unsaturated ketone, using sodium hypochlorite, is illustrated in Scheme 1.8 [10]. Hypochlorite is extracted as an onium


Starks Extraction Mechanism



Brändström–Montanari Modification



Scheme 1.7

Asymmetric Epoxidation of α,β -Unsaturated Ketones

Scheme 1.8

salt into the organic phase, and the onium hypochlorite then reacts with the ketone while recognizing its enantiotopic face to provide the optically enriched α,β -epoxy ketone.

1.4

Use of Crown Ether as Phase-Transfer Catalyst

Crown ether is the other important class of phase-transfer catalysts which critically differs from the onium salt, in that the whole inorganic salt is transferred into the organic phase. The reaction modes described above can generally be accommodated in such crown ether-catalyzed reactions, simply by replacing the onium cation by a metal cation complex of crown ether.

References

- 1 Starks, C.M. (1971) *J. Am. Chem. Soc.*, **93**, 195.
- 2 Makosza's pioneering contributions are representative, see: (a) Makosza, M. (1966) *Tetrahedron Lett.*, 4621; (b) Makosza, M. (1966) *Tetrahedron Lett.*, 5489; (c) Makosza, M. (1969) *Tetrahedron Lett.*, 673; (d) Makosza, M. (1969) *Tetrahedron Lett.*, 677; see also: (e) Brändström, A. (1977) *Adv. Phys. Org. Chem.*, **15**, 267.
- 3 (a) Dehmlow, E.V. and Dehmlow, S.S. (1993) *Phase Transfer Catalysis*, 3rd edn. VCH, Weinheim; (b) Starks, C.M., Liotta, C.L. and Halpern, M.E. (1994) *Phase-Transfer Catalysis*, Chapman & Hall, New York; (c) Sasson, Y. and Neumann, R. (eds), (1997) *Handbook of Phase-Transfer Catalysis*, Blackie Academic & Professional, London; (d) Halpern, M.E. (ed.), (1997) *Phase-Transfer Catalysis*, ACS