

Springer Climate

Małgorzata Falarz *Editor*


Climate Change in Poland

Past, Present, Future

 Springer

Springer Climate

Series Editor

John Dodson , Institute of Earth Environment, Chinese Academy of Sciences,
Xian, Shaanxi, China

Springer Climate is an interdisciplinary book series dedicated to climate research. This includes climatology, climate change impacts, climate change management, climate change policy, regional climate studies, climate monitoring and modeling, palaeoclimatology etc. The series publishes high quality research for scientists, researchers, students and policy makers. An author/editor questionnaire, instructions for authors and a book proposal form can be obtained from the Publishing Editor.

Now indexed in Scopus® !

More information about this series at <http://www.springer.com/series/11741>

Małgorzata Falarz
Editor

Climate Change in Poland

Past, Present, Future

 Springer

Editor

Małgorzata Falarz 
Faculty of Natural Sciences
Institute of Earth Sciences
University of Silesia in Katowice
Katowice, Poland

ISSN 2352-0698

ISSN 2352-0701 (electronic)

Springer Climate

ISBN 978-3-030-70327-1

ISBN 978-3-030-70328-8 (eBook)

<https://doi.org/10.1007/978-3-030-70328-8>

© Springer Nature Switzerland AG 2021, corrected publication 2021

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, expressed or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Switzerland AG
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

To our Masters

Authors

Preface

This monograph would not have been created without the activity and determination of its authors, 36 Polish climatologists who contributed its release. Authorship of individual book chapters was entrusted to the experienced scientists, specialists in the research of individual elements and characteristics of climate. The authors of the monograph come from eight scientific centres in Poland (Gdańsk, Katowice/Sosnowiec, Kraków, Łódź, Lublin, Poznań, Toruń, Warszawa), representing eight universities (Adam Mickiewicz University in Poznań, Jagiellonian University in Kraków, Maria Curie-Skłodowska University in Lublin, Nicolaus Copernicus University in Toruń, the University of Gdańsk, the University of Łódź, the University of Silesia in Katowice, the University of Warsaw), two technical universities (Kraków University of Technology, Agricultural University of Kraków), the Polish Academy of Science and Institute of Meteorology and Water Management—National Research Institute. The vast majority of the research presented here is new and was conducted for the purposes of this monograph. However, the book was preceded by many works, publications and studies that influenced its form.

This book would not have been created without the tireless work of many unnamed observers at the weather stations (both the Institute of Meteorology and Water Management—National Research Institute and university stations), who, day after day, have painstakingly measured, observed and documented the state of the weather for many years. In addition, the book could not have been published without the creation and availability of a meteorological database by the Institute of Meteorology and Water Management—National Research Institute and universities.

Finally, the monograph would not have been published without the active help of English native speaker, Leah Morawiec, who took care of the linguistic correctness of most chapters of the book. Special thanks are due to the scientific and technical staff of Springer International Publishing, especially for their great patience and kind help.

I express my deep gratitude to everyone for their work, commitment and effort.

The book is addressed to scientists (climatologists, geographers, etc.), academic teachers, students, journalists, and all those interested in Poland and climate change in Poland.

Climate variability and its contemporary changes are happening right in front of us. Therefore, we would appreciate any feedback to help improve the next editions of the book.

Sosnowiec, Poland

Małgorzata Falarz

Acknowledgments In this monograph meteorological data of Institute of Meteorology and Water Management—National Research Institute and meteorological stations of both Jagiellonian University in Kraków (stations: Kraków Obs. UJ and Gaik-Brzezowa) and University of Silesia in Katowice (station Sosnowiec) has been used and processed.

The original version of the book was revised: In Chapter 11 figures 23 and 33 corrections have been updated and also the affiliation city name “Katowice” of the volume editor is corrected in web version. The correction to the book is available at https://doi.org/10.1007/978-3-030-70328-8_25

Contents

Part I Background

- 1 Introduction** 3
Małgorzata Falarz, Rajmund Przybylak, Janusz Filipiak,
Agnieszka Wypych, and Małgorzata Szwed
- 2 Initial Research of Climate Change in Poland** 9
Małgorzata Falarz, Magdalena Opała-Owczarek,
Tadeusz Niedźwiedź, Zuzanna Bielec-Bąkowska,
Jakub Wojkowski, Dorota Matuszko, Janusz Filipiak,
Zbigniew Ustrnul, Agnieszka Wypych, Monika Lisowska,
Anita Bokwa, Krzysztof Błażejczyk, Katarzyna Piotrowicz,
and Małgorzata Szwed
- 3 Data and Methods of Investigation** 29
Małgorzata Falarz, Małgorzata Szwed,
Magdalena Opała-Owczarek, Magdalena Kuchcik,
Łukasz Małarzewski, Katarzyna Piotrowicz,
Tadeusz Niedźwiedź, Zuzanna Bielec-Bąkowska, Anita Bokwa,
Janusz Filipiak, Dorota Matuszko, Zbigniew Ustrnul,
Joanna Wibig, and Agnieszka Wypych
- 4 Homogeneity of Climate Series** 45
Małgorzata Falarz, Joanna Wibig, Dorota Matuszko,
Janusz Filipiak, Monika J. Hajto, Mateusz Taszarek,
Zuzanna Bielec-Bąkowska, Agnieszka Wypych, and Zbigniew Ustrnul

Part II Long-Term Climate Change

- 5 Climate Change Before Instrumental Measurements** 71
Magdalena Opała-Owczarek, Tadeusz Niedźwiedź,
Rajmund Przybylak, and Wojciech Tylmann

Part III Recent Climate Change

6	Change of Atmospheric Circulation	123
	Tadeusz Niedźwiedź and Zbigniew Ustrnul	
7	Air Pressure Change	151
	Zuzanna Bielec-Bąkowska and Katarzyna Piotrowicz	
8	Solar Radiation Change	177
	Małgorzata Falarz, Łukasz Małarzewski, Joanna Uscka-Kowalkowska, Dorota Matuszko, and Tomasz Budzik	
9	Change of Sunshine	189
	Dorota Matuszko, Stanisław Węglarczyk, Krzysztof Bartoszek, Jakub Soroka, and Barbara Bogdańska	
10	Change of Cloudiness	217
	Janusz Filipiak	
11	Air Temperature Change	275
	Zbigniew Ustrnul, Agnieszka Wypych, and Danuta Czekierda	
12	Air Humidity Change	331
	Agnieszka Wypych	
13	Precipitation Change	349
	Ewa Łupikasza and Łukasz Małarzewski	
14	Snow Cover Change	375
	Małgorzata Falarz and Ewa Bednorz	
15	Change of Wind	391
	Joanna Wibig	
16	Change of Thunderstorms and Tornadoes	421
	Zuzanna Bielec-Bąkowska, Mateusz Taszarek, and Leszek Kolendowicz	
17	Change of Hail Frequency	443
	Zuzanna Bielec-Bąkowska	
18	Change of Fog Frequency	453
	Anita Bokwa, Agnieszka Wypych, and Monika J. Hajto	
19	Changes in Bioclimatic Indices	471
	Magdalena Kuchcik, Krzysztof Błażejczyk, and Agnieszka Halaś	
20	Change of Weather Types	493
	Katarzyna Piotrowicz and Dominika Ciaranek	

Part IV Future Climate Change

- 21 Projections of Temperature Changes in Poland** 513
Małgorzata Szwed
- 22 Projections of Precipitation Changes in Poland** 529
Iwona Pińskwar and Adam Choryński
- 23 Projected Changes in Thermal Indices Related
to the Agriculture and Energy Sectors** 545
Dariusz Graczyk, Iwona Pińskwar, and Adam Choryński

Part V Discussion and Conclusion

- 24 Climate Change in Poland—Summary, Discussion
and Conclusion** 561
Małgorzata Falarz, Zuzanna Bielec-Bąkowska,
Agnieszka Wypych, Dorota Matuszko, Tadeusz Niedźwiedź,
Iwona Pińskwar, Katarzyna Piotrowicz, Zbigniew Ustrnul,
Anita Bokwa, Janusz Filipiak, Dariusz Graczyk,
Magdalena Kuchcik, Magdalena Opała-Owczarek,
Małgorzata Szwed, and Joanna Wibig
- Correction to: Climate Change in Poland** C1
Małgorzata Falarz

Contributors

Krzysztof Bartoszek Faculty of Earth Sciences and Spatial Management, Institute of Earth and Environmental Sciences, Department of Hydrology and Climatology, Maria Curie-Skłodowska University, Lublin, Poland

Ewa Bednorz Faculty of Geographical and Geological Sciences, Institute of Physical Geography and Environmental Planning, Department of Meteorology and Climatology, Adam Mickiewicz University, Poznań, Poland

Zuzanna Bielec-Bąkowska Faculty of Natural Sciences, Institute of Earth Sciences, University of Silesia in Katowice, Katowice, Poland

Krzysztof Błażejczyk Institute of Geography and Spatial Organization, Past Landscape Dynamic Laboratory, Polish Academy of Sciences, Warsaw, Poland

Barbara Bogdańska Institute of Meteorology and Water Management—National Research Institute, Warsaw, Poland

Anita Bokwa Faculty of Geography and Geology, Institute of Geography and Spatial Management, Jagiellonian University, Kraków, Poland

Tomasz Budzik Faculty of Natural Sciences, Institute of Earth Sciences, University of Silesia in Katowice, Katowice, Poland

Adam Choryński Institute for Agricultural and Forest Environment, Polish Academy of Sciences, Poznań, Poland

Dominika Ciaranek Faculty of Geography and Geology, Institute of Geography and Spatial Management, Jagiellonian University, Kraków, Poland

Danuta Czekierda Institute of Meteorology and Water Management—National Research Institute, Warsaw, Poland

Małgorzata Falarz Faculty of Natural Sciences, Institute of Earth Sciences, University of Silesia in Katowice, Katowice, Poland

Janusz Filipiak Faculty of Oceanography and Geography, Institute of Geography, Department of Meteorology and Climatology, University of Gdańsk, Gdańsk, Poland;

Institute of Meteorology and Water Management—National Research Institute, Warsaw, Poland

Dariusz Graczyk Institute for Agricultural and Forest Environment, Polish Academy of Sciences, Poznań, Poland

Monika J. Hajto Institute of Meteorology and Water Management—National Research Institute, Warsaw, Poland;

Faculty of Geography and Geology, Institute of Geography and Spatial Management, Jagiellonian University, Kraków, Poland

Agnieszka Halaś Institute of Geography and Spatial Organization, Past Landscape Dynamic Laboratory, Polish Academy of Sciences, Warsaw, Poland

Leszek Kolendowicz Faculty of Geographical and Geological Sciences, Institute of Physical Geography and Environmental Planning, Department of Meteorology and Climatology, Adam Mickiewicz University, Poznań, Poland

Magdalena Kuchcik Institute of Geography and Spatial Organization, Climate Impacts Laboratory, Polish Academy of Sciences, Warsaw, Poland

Monika Lisowska Faculty of Geography and Regional Studies, Chair of Physical Geography, Department of Climatology, University of Warsaw, Warsaw, Poland

Ewa Łupikasza Faculty of Natural Sciences, Institute of Earth Sciences, University of Silesia in Katowice, Katowice, Poland

Łukasz Małarzewski Faculty of Natural Sciences, Institute of Earth Sciences, University of Silesia in Katowice, Katowice, Poland

Dorota Matuszko Faculty of Geography and Geology, Institute of Geography and Spatial Management, Jagiellonian University in Kraków, Kraków, Poland

Tadeusz Niedźwiedź Faculty of Natural Sciences, Institute of Earth Sciences, University of Silesia in Katowice, Katowice, Poland

Magdalena Opała-Owczarek Faculty of Natural Sciences, Institute of Earth Sciences, University of Silesia in Katowice, Katowice, Poland

Iwona Pińskwar Institute for Agricultural and Forest Environment, Polish Academy of Sciences, Poznań, Poland

Katarzyna Piotrowicz Faculty of Geography and Geology, Institute of Geography and Spatial Management, Jagiellonian University, Kraków, Poland

Rajmund Przybylak Faculty of Earth Sciences and Spatial Management, Department of Meteorology and Climatology, Nicolaus Copernicus University, Toruń, Poland;

Centre for Climate Change Research, Nicolaus Copernicus University, Toruń, Poland

Jakub Soroka Institute of Meteorology and Water Management—National Research Institute, Hydrological and Meteorological Station, Gorzów Wielkopolski, Poland

Małgorzata Szwed Institute for Agricultural and Forest Environment, Polish Academy of Sciences, Poznań, Poland

Mateusz Taszarek Faculty of Geographical and Geological Sciences, Institute of Physical Geography and Environmental Planning, Department of Meteorology and Climatology, Adam Mickiewicz University, Poznań, Poland

Wojciech Tylmann Faculty of Oceanography and Geography, Environmental Change Reconstruction Laboratory, University of Gdansk, Gdańsk, Poland

Joanna Uscka-Kowalkowska Faculty of Earth Sciences and Spatial Management, Department of Meteorology and Climatology, Nicolaus Copernicus University, Toruń, Poland

Zbigniew Ustrnul Faculty of Geography and Geology, Institute of Geography and Spatial Management, Jagiellonian University, Kraków, Poland;
Institute of Meteorology and Water Management—National Research Institute, Warsaw, Poland

Stanisław Węglarczyk Faculty of Environmental and Power Engineering, Department of Geoengineering and Water Management, Cracow University of Technology, Kraków, Poland

Joanna Wibig Faculty of Geographical Sciences, Department of Meteorology and Climatology, University of Lodz, Lodz, Poland

Jakub Wojkowski Faculty of Environmental Engineering and Geodesy, Department of Ecology, Climatology and Air Protection, University of Agriculture, Kraków, Poland

Agnieszka Wypych Faculty of Geography and Geology, Institute of Geography and Spatial Management, Jagiellonian University in Kraków, Kraków, Poland;
Institute of Meteorology and Water Management—National Research Institute, Warsaw, Poland

Part I

Background

Chapter 1

Introduction



**Małgorzata Falarz, Rajmund Przybylak, Janusz Filipiak,
Agnieszka Wypych, and Małgorzata Szwed**

Abstract Climate change is now one of the most important global challenges. The problem was noticed worldwide as early as the 1970s, when the first world Climate Conference was held in Stockholm (1972) under the motto “Only One Earth”. Great progress, however, did not begin in climate protection until 1992, when the Rio de Janeiro Earth Summit was organised. Later on, since 1995, the Climate Summits (Conferences of the Parties, COP) of the United Nations have been held every year, bringing together representatives from almost all countries of the world and resulting in important global agreements on combating global warming (e.g. the Kyoto Protocol in 1997, Paris Agreement in 2015).

M. Falarz (✉)

Faculty of Natural Sciences, Institute of Earth Sciences, University of Silesia in Katowice,
Katowice, Poland

e-mail: malgorzata.falarz@us.edu.pl

R. Przybylak

Faculty of Earth Sciences and Spatial Management, Department of Meteorology and Climatology,
Nicolaus Copernicus University, Toruń, Poland

Centre for Climate Change Research, Nicolaus Copernicus University, Toruń, Poland

R. Przybylak

e-mail: rp11@umk.pl

J. Filipiak

Faculty of Oceanography and Geography, Institute of Geography, Department of Meteorology and
Climatology, University of Gdańsk, Gdańsk, Poland

e-mail: janusz.filipiak@ug.edu.pl

J. Filipiak · A. Wypych

Institute of Meteorology and Water Management—National Research Institute, Warsaw, Poland

e-mail: agnieszka.wypych@uj.edu.pl

A. Wypych

Faculty of Geography and Geology, Institute of Geography and Spatial Management, Jagiellonian
University, Krakow, Poland

M. Szwed

Institute for Agricultural and Forest Environment, Polish Academy of Sciences, Poznań, Poland

e-mail: mszwed@man.poznan.pl

© Springer Nature Switzerland AG 2021

M. Falarz (ed.), *Climate Change in Poland*, Springer Climate,

https://doi.org/10.1007/978-3-030-70328-8_1

Climate change is now one of the most important global challenges. The problem was noticed worldwide as early as the 1970s, when the first world Climate Conference was held in Stockholm (1972) under the motto “Only One Earth”. Great progress, however, did not begin in climate protection until 1992, when the Rio de Janeiro Earth Summit was organised. Later on, since 1995, the Climate Summits (Conferences of the Parties, COP) of the United Nations have been held every year, bringing together representatives from almost all countries of the world and resulting in important global agreements on combating global warming (e.g. the Kyoto Protocol in 1997, Paris Agreement in 2015).

The Intergovernmental Panel on Climate Change (IPCC), the United Nations body for assessing the science related to climate change, publishes reports every 5–7 years with scientific assessments on climate change (e.g. IPCC 1992, 2007, 2013), its implications and potential future risks. In addition, special reports are published. The report appearing in 2018 for example, highlights the uneven distribution of temperature values on the globe for scenarios of 1.5 °C and 2.0 °C global warming compared to the preindustrial level. In line with expectation, the biggest changes will be in the extreme temperature in land areas, e.g. with global mean surface temperature warming of 2.0 °C, the change in average temperature of hottest days reaches 3–4 °C in some land areas, including Europe (IPCC 2018).

A number of climate change studies have been carried out worldwide, the results of which have been published in thousands of scientific articles or multifaceted monographs (e.g. Jones et al. 1996, Houghton et al. 2001, Jones et al. 2001, Hurrell et al. 2003, Knieling and Leal Filho 2013, Leal Filho et al. 2016).

Interest in climate change in Poland has a long history and tradition, for review see e.g. Przybylak et al. 2010 and Section I.2 in this volume. Here we mention only some of the more recent and most important works. They generally cover many aspects of climate change in Poland, including analyses of: (1) climate before instrumental measurements (e.g. Przybylak et al. 2010; Opała 2015; Przybylak 2016), and (2) contemporary climate: circulation conditions (e.g. Niedźwiedz and Łupikasza 2016, 2019), air pressure (e.g. Bielec-Bąkowska 2016), solar radiation (e.g. Uscka-Kowalkowska et al. 2007), sunshine (e.g. Matuszko 2014; Matuszko et al. 2020), cloudiness (e.g. Filipiak and Miętus 2009), air temperature (e.g. Ustrnul 2000), air humidity (e.g. Wypych 2010), precipitation (e.g. Łupikasza 2010, 2016), snow cover (e.g. Falarz 2008), wind (e.g. Arażny et al. 2007; Lorenc 2012), thunderstorms (e.g. Bielec-Bąkowska 2003) and tornadoes (e.g. Taszarek and Brooks 2015), hail (e.g. Bielec-Bąkowska 2013), fog (e.g. Łupikasza and Niedźwiedz 2016), urban climate (Fortuniak 2019), bioclimatic indices (e.g. Błażejczyk and Matzarakis 2007), weather types (e.g. Piotrowicz et al. 2016), temperature projections (e.g. Szwed and Graczyk 2006), precipitation projections (e.g. Pińskwar 2010) and projected changes in indices related to agriculture, water resources and human health (e.g. Szwed et al. 2010).

In the last 20 years there have been several conference publications in Polish presenting: circulation and radiation conditions of climate change and variability in Poland (Kolendowicz 2010; Bielec-Bąkowska et al. 2012; Kolendowicz et al. 2019), climate change in Poland at different spatial and time scales (Piotrowicz

and Twardosz 2007), thermal and precipitation variability (Bednorz 2010), thermal, radiation and hydrological aspects of change (Chojnacka-Ożga and Lorenc 2019) and climate change and its consequences with references to the climate of Europe (Zmiany i zmienność klimatu Polski [...] 1999); Bednorz and Kolendowicz 2010).

Particular attention has been paid to extreme meteorological and hydrological phenomena in Poland (Kundzewicz and Jania 2007; Ustrnul and Czekierda 2009).

Of particular note are monographs in English, with an international composition of authors presenting: climate change in Poland in relation to European change (Przybylak et al. 2010) and climate change of the Baltic Sea Basin area covering Poland (The BACC Author Team 2008; The BACC II Author Team 2015). The first (Przybylak et al. 2010) provides both a synthesis of current knowledge about climate of Poland in recent centuries and new finding results for selected issues in selected regions of Poland or for individual meteorological stations. The book relates to the results published till 2009.

The above-mentioned works were created in different years, using different methods, with different levels of detail. None of them undertakes a systematic spatial description of the change of all climate elements throughout the area of Poland. The knowledge of Poland's climate change so far is very rich and valuable. However, it is somewhat dispersed; there is no harmonisation of methods, databases or research periods. There is not yet a book about climate change in Poland which presents new research that uses an uniform method for all the elements of climate as well as for the selected meteorological phenomena, bioclimatic conditions and weather types for past, present and future change. In this book, knowledge of these subjects is sorted out and harmonised.

The current volume consists of five main parts. Part I is an introduction. Part II presents the results of the study of climate change before the beginning of the instrumental measurements in Poland in the last millennium. In Part III the long-term trends and variability of circulation indices, climate elements, selected meteorological phenomena, bioclimatic conditions and weather types are analysed using data from the Institute of Meteorology and Water Management–National Research Institute and two university stations. A spatial diversity of climate trends and variability in Poland is presented in maps. Part IV of the book deals with projected changes up to 2100 in temperature, precipitation and thermal indices related to the agriculture and energy sectors in Poland. Part V contains summary, discussion and conclusion. More than 80% of the results presented in this monograph are the results of new studies, not yet published, which were carried out specifically for the purposes of this book.

References

- Araźny A, Przybylak R, Vizi Z et al (2007) Mean and extreme wind velocities in Central Europe 1951–2005. *Geogr Pol* 80(2):69–78

- Bednorz E (ed) (2010) *Klimat Polski na tle klimatu Europy. Warunki termiczne i opadowe*, Studia i Prace z Geografii i Geologii, 15, Bogucki Wydawnictwo Naukowe, Poznań, 214 pp
- Bednorz E, Kolendowicz L (eds) (2010) *Klimat Polski na tle klimatu Europy. Zmiany i ich konsekwencje*, Studia i Prace z Geografii i Geologii, 16, Bogucki Wydawnictwo Naukowe, Poznań, 244 pp
- Bielec-Bąkowska Z (2003) Long-term variability of thunderstorms occurrence in Poland in the 20th century. *Atmos Res* 67–68:35–52
- Bielec-Bąkowska Z (2013) Burze i grady w Polsce. *Prace Geograficzne Instytutu Geografii i Gospodarki Przestrzennej UJ* 132:99–132
- Bielec-Bąkowska Z (2016) Long-term variability of the frequency and persistence of strong highs over Poland. *Environ Socio-Econ Stud* 4(1):12–23
- Bielec-Bąkowska Z, Łupikasza E, Widawski A (eds) (2012) Rola cyrkulacji atmosfery w kształtowaniu klimatu; *Prace Wydziału Nauk o Ziemi Uniwersytetu Śląskiego*, 74, Sosnowiec
- Błażejczyk K, Matzarakis A (2007) Assessment of bioclimatic differentiation of Poland based on the human heat balance. *Geogr Pol* 80(1):63–82
- Chojnacka-Ożga L, Lorenc H (eds) (2019) *Współczesne problemy klimatu Polski*, Institute of Meteorology and Water Management—National Research Institute, Warsaw, Poland, 260 pp
- Falarz M (2008) Changes of extreme nival conditions in Poland during the second half of the 20th century. *Meteorol Z* 17(3):339–444
- Filipiak J, Miętus M (2009) Spatial and temporal variability of cloudiness in Poland, 1971–2000. *Int J Climatol* 29:1294–1311. <https://doi.org/10.1002/joc.1777>
- Fortuniak K (2019) Studies on urban climate in Poland. *Przegląd Geofizyczny LXIV* 1–2:73–106. <https://doi.org/10.32045/PG-2019-003>
- Houghton JT et al (eds) (2001) *Climate change 2001: the scientific basis*. Cambridge University Press, Cambridge, UK
- Hurrell JW, Kushnir Y, Ottensen G, Visbeck M (eds) (2003) *North Atlantic oscillation: climatic significance and environmental impact*. Geophysical Monograph 134. American Geophysical Union, Washington
- IPCC (1992) *Climate change: the 1990 and 1992 IPCC Assessments Report*. JT Houghton (ed), Cambridge, UK, 212 pp
- IPCC (2007) *Climate change 2007, The Physical Science Basis. Contribution of working group I to the fourth assessment report of the intergovernmental panel on climate change*. In: Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 996 pp
- IPCC (2013) *Climate change 2013, The physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change*. In: Stocker TF, Qin D, Plattner G.-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM (eds). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp. <https://doi.org/10.1017/cbo9781107415324>
- IPCC (2018) *Global Warming of 1.5 °C. An IPCC Special Report on the impacts of global warming of 1.5 °C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty*. In: Masson-Delmotte V, Zhai P, Pörtner H-O, Roberts D, Skea J, Shukla PR, Pirani A, Moufouma-Okia W, Péan C, Pidcock R, Connors S, Matthews JBR, Chen Y, Zhou X, Gomis MI, Lonnoy E, Maycock T, Tignor M, Waterfield T (eds)
- Jones PD, Bradley RS, Jouzel J (eds) (1996) *Climatic variations and forcing mechanisms of the last 2000 years*. Springer, Berlin/Heidelberg/New York
- Jones PD, Ogilvie AEJ, Davies TD, Briffa KR (eds) (2001) *History and climate, memories of the future*. Kluwer, New York
- Knieling J, Leal Filho W (eds) (2013) *Climate change governance*. Springer International Publishing, 320 pp
- Kolendowicz L (ed) (2010) *Klimat Polski na tle klimatu Europy. Warunki cyrkulacyjne i radiacyjne*, Studia i Prace z Geografii i Geologii, 14, Bogucki Wydawnictwo Naukowe, Poznań, 170 pp

- Kolendowicz L, Bednorz E, Tomczyk AM (2019) Zmienność klimatu Polski i Europy oraz jej cyrkulacyjne uwarunkowania, *Studia i Prace z Geografii*, 77, Bogucki Wydawnictwo Naukowe, Poznań, 264 pp
- Kundzewicz ZW, Jania J (2007) Extreme meteorological and hydrological events in Poland. *Geogr Pol*, 80(2)
- Leal Filho W, Musa H, Cavan G, O'Hare P, Seixas J (eds) (2016) *Climate change adaptation*. Springer International Publishing, Resilience and Hazards, p 459
- Lorenc H (2012) Maksymalne prędkości wiatru w Polsce. IMGW PIB, Warszawa
- Łupikasza E (2010) Spatial and temporal variability of extreme precipitation in Poland in the period 1951–2006. *Int J Climatol* 30:991–1007. <https://doi.org/10.1002/joc.1950>
- Łupikasza E (2016) *The climatology of air-mass and frontal extreme precipitation. Study of meteorological data in Europe*. Springer International Publishing, 313 pp
- Łupikasza E, Niedźwiedz T (2016) Synoptic climatology of fog in selected locations of southern Poland (1966–2015). *Bull Geogr, Phys Geogr Ser* 11:5–15
- Matuszko D (2014) Long-term variability in solar radiation in Krakow based on measurements of sunshine duration. *Int J Climatol* 34:228–234
- Matuszko D, Bartoszek K, Soroka J, Węglarczyk S (2020) Sunshine duration in Poland from ground- and satellite-based data. *Int J Climatol*. <https://doi.org/10.1002/joc.6460>
- Niedźwiedz T, Łupikasza E (2016) Change in atmospheric circulation patterns In: Kundzewicz ZW, Stoffel M, Niedźwiedz T, Wyźga B (eds) *Flood risk in the Upper Vistula Basin*. Springer International Publishing Switzerland, pp 189–208, https://doi.org/10.1007/978-3-319-41923-7_9
- Niedźwiedz T, Łupikasza E (2019) Atmospheric circulation in the investigations of Polish climatologists. *Przegląd Geofizyczny LXIV* 1–2:107–166. <https://doi.org/10.32045/PG-2019-004>
- Opała M (2015) The 443-year tree-ring chronology for the scots pine from Upper Silesia (Poland) as a dating tool and climate proxy. *Geochronometria* 42(1):41–52
- Pińskwar I (2010) Projekcje zmian w ekstremach opadowych w Polsce. (Projections of changes in extreme precipitation in Poland) *Monografia KGW PAN*, 34 (in Polish)
- Piotrowicz K, Bielec-Bąkowska Z, Domonkos P (2016) High atmospheric pressure and accompanying cold season weather types in Poland (1951 – 2010). *Clim Res* 67:165–177. <https://doi.org/10.3354/cr01364>
- Piotrowicz K, Twardosz R (eds) (2007) *Wahania klimatu w różnych skalach przestrzennych i czasowych*. IGI GP UJ, Kraków, p 428
- Przybylak R (2016) Poland's climate in the Last Millennium, *Oxford Research Encyclopedia, Climate Science*, Oxford University Press USA, 36 pp., <https://doi.org/10.1093/acrefore/9780190228620.013.2>
- Przybylak R, Majorowicz J, Brázdil R, Kejna M (Eds.) (2010) The Polish climate in the European context: an historical overview, Springer, 535 pp. <https://doi.org/10.1007/978-90-481-3167-9>
- Szwed M, Graczyk D (2006) Thermal seasons in Poland—the present and the future based on HadRM3-PRECIS results. *Pap Glob Chang* 13:59–77
- Szwed M, Karg G, Pińskwar I, Radziejewski M, Graczyk D, Kędziora A, Kundzewicz ZW (2010) Climate change and its effect on agriculture, water resources and human health sectors in Poland. *Nat Hazards Earth Syst Sci* 10:1725–1737. <https://doi.org/10.5194/nhess-10-1725-2010>
- Taszarek M, Brooks HE (2015) Tornado climatology of Poland. *Mon Weather Rev* 143(3):702–717
- The BACC Author Team (2008) *Assessment of climate change for the Baltic Sea Basin*. Springer-Verlag, Berlin Heidelberg
- The BACC II Author Team (2015) *Second assessment of climate change for the Baltic Sea Basin*, ser.: *Regional Climate Studies*, Springer, London
- Uscka-Kowalkowska J, Przybylak R, Vízi Z, Araźny A, Kejna M, Maszewski R (2007) Variability to global solar radiation in Central Europe during the period 1951–2005 (on the basis of data from NCEP/NCAR reanalysis project). *Geogr Pol* 80(2):59–68
- Ustrnul Z (2000) Synoptic-climatic structure of the extreme air thermal phenomena in Poland. *Geogr Pol* 73(2):99–109

- Ustrnul Z, Czekierda D (2009) Atlas ekstremalnych zjawisk meteorologicznych oraz sytuacji synoptycznych w Polsce. IMGW, Warszawa
- Wypych A (2010) Twentieth century variability of surface humidity as the climate change indicator in Kraków (Southern Poland). *Theor Appl Climatol* 101:475–482
- Zmiany i zmienność klimatu Polski. Ich wpływ na gospodarkę, ekosystemy i człowieka (1999) Ogólnopolska konferencja naukowa, Łódź, 4–6 listopada 1999 r., 315 pp

Chapter 2

Initial Research of Climate Change in Poland



Małgorzata Falarz, Magdalena Opała-Owczarek, Tadeusz Niedźwiedź, Zuzanna Bielec-Bąkowska, Jakub Wojkowski, Dorota Matuszko, Janusz Filipiak, Zbigniew Ustrnul, Agnieszka Wypych, Monika Lisowska, Anita Bokwa, Krzysztof Błażejczyk, Katarzyna Piotrowicz, and Małgorzata Szwed

Abstract This chapter includes the results of the earliest studies on changes and variability of climatic elements, bioclimatic indices and weather types across Poland. The first pioneering works on climatic studies were presented (since 1858), even if they did not relate to climate change.

M. Falarz (✉) · M. Opała-Owczarek · T. Niedźwiedź · Z. Bielec-Bąkowska
Faculty of Natural Sciences, Institute of Earth Sciences, University of Silesia in Katowice,
Katowice, Poland
e-mail: malgorzata.falarz@us.edu.pl

M. Opała-Owczarek
e-mail: magdalena.opala@us.edu.pl

T. Niedźwiedź
e-mail: tadeusz.niedzwiedz@us.edu.pl

Z. Bielec-Bąkowska
e-mail: zuzanna.bielec-bakowska@us.edu.pl

J. Wojkowski
Faculty of Environmental Engineering and Geodesy, Department of Ecology, Climatology and Air
Protection, University of Agriculture, Kraków, Poland
e-mail: jakub.wojkowski@urk.edu.pl

D. Matuszko · Z. Ustrnul · A. Wypych · A. Bokwa · K. Piotrowicz
Faculty of Geography and Geology, Institute of Geography and Spatial Management, Jagiellonian
University, Kraków, Poland
e-mail: d.matuszko@uj.edu.pl

Z. Ustrnul
e-mail: zbigniew.ustrnul@uj.edu.pl

A. Wypych
e-mail: agnieszka.wypych@uj.edu.pl

A. Bokwa
e-mail: anita.bokwa@uj.edu.pl

K. Piotrowicz
e-mail: k.piotrowicz@uj.edu.pl

This chapter includes the results of the earliest studies on changes and variability of climatic elements, bioclimatic indices and weather types across Poland. The first pioneering works on climatic studies were presented (since 1858), even if they did not relate to climate change. Already in the nineteenth century, the results of air temperature (e.g. Karliński 1868; Kuczyński 1884), precipitation (e.g. Kremser 1884; Wild 1887), cloudiness (Satke 1898), humidity (Wierzbicki 1878) and snow cover (Satke 1896, 1899) investigations in Polish lands were published in Polish and German.

The climatic history in Poland for the **preinstrumental period** has been reconstructed from various natural archives. Information on climate change during the Holocene period was obtained on the basis of geomorphological, sedimentological and botanical data (Stasiak 1968; Starkel 1977; Ralska-Jasiewiczowa and Starkel 1988, 1991; Starkel et al. 1996; Kotarba and Baumgart-Kotarba 1997) and the most accurate climatic reconstructions were carried out for the historical period (the last millennium) (Niedźwiedź et al. 2015). Based on a literature review, it can be concluded that information on climate change in Poland during historical times were provided mainly by four types of proxy data: documentary evidences, tree rings, lake sediments and geothermal profiles.

The earliest study on Poland's climatic conditions was conducted using documentary evidences (Semkowicz 1922; Polaczkówna 1925). A considerable amount of data can be found in the excerpts from the chronicles (e.g. Walawender 1932; Namaczyńska 1937; Szewczuk 1939; Inglot 1962, 1966, 1968, Rojecki 1965); however, the first climatological interpretation of these historical records was made no longer than 30 years ago by Maruszczak (1988), Sadowski (1991), Limanówka (2001), and Przybylak et al. (2001). Detailed information on the development of historical climatology in Poland was presented by Przybylak et al. (2001).

J. Filipiak

Faculty of Oceanography and Geography, Institute of Geography, Department of Meteorology and Climatology, University of Gdańsk, Gdańsk, Poland
e-mail: janusz.filipiak@ug.edu.pl

J. Filipiak · Z. Ustrnul · A. Wypych

Institute of Meteorology and Water Management—National Research Institute, Warsaw, Poland

M. Lisowska

Faculty of Geography and Regional Studies, Chair of Physical Geography, Department of Climatology, University of Warsaw, Warsaw, Poland
e-mail: mlisowska@uw.edu.pl

K. Błażejczyk

Polish Academy of Sciences, Institute of Geography and Spatial Organization, Climate Impacts Laboratory, Warsaw, Poland
e-mail: kblaz@twarda.pan.pl

M. Szwed

Institute for Agricultural and Forest Environment, Polish Academy of Sciences, Poznań, Poland
e-mail: mszwed@man.poznan.pl

The earliest mention of the possibility of conducting dendroclimatological research in Poland comes from 1914 (Merecki 1914). Although preliminary dendroclimatological research started after the Second World War (Zinkiewicz 1946; Ermich 1953), the first reconstruction of climate on the basis of tree rings was presented by Bednarz (1984) for the Tatra Mountains. Since these pioneer work many long tree-ring chronologies were developed for different parts of the country (see summary presented by Zielski et al. 2010). However, development of studies on past climatic changes in Poland based on dendroclimatic records has taken place in the last two decades (e.g. Krąpiec et al. 1998; Cedro 2004; Niedźwiedz 2004; Büntgen et al. 2007; Szychowska-Krąpiec 2010; Koprowski et al. 2012; Opała and Mendecki 2014; Balanzategui et al. 2017).

Another source of paleoclimatic data covering the last millennium is laminated lake sediments, whose potential for paleoenvironmental and paleoclimatic reconstructions has been thoroughly examined for the Lake Gościąg site in central Poland (Ralska-Jasiewiczowa et al. 1992, 1998; Starkel et al. 1996). Recently, high potential for quantitative paleoenvironmental reconstructions was also shown for Lake Żabińskie in north-eastern Poland (Amann et al. 2014; Hernández-Almeida et al. 2015; Larocque-Tobler et al. 2015).

The first use of geothermal profiles in thermally stabilised wells to reconstruct temperature history in Poland over the last 500 years was made by Majorowicz et al. (2001).

In the first half of the twentieth century, there was no work on the long-term variability of **atmospheric circulation** over Poland, but there were publications on the impact of atmospheric circulation on some elements of the climate (Bartnicki 1924; Arctowski 1927; Kaczorowska 1933; Milata 1935; Bartnicki and Kołodziejczyk 1935; Lisowski 1935). In some later works, the role of circulation factors in the formation of the Polish climate was emphasised (Kozuchowski 2003, 2004; Miętus 1996; Wibig 2001). In the second half of the twentieth century, several classifications of circulation types and indices were created (Osuchowska-Klein 1975; Niedźwiedz 1988; Ustrnul 1997; Piotrowski 2009). Attention was also paid to the long-term variability of circulation types, air masses and atmospheric fronts (Parczewski 1964; Niedźwiedz 1996; Kaszewski and Filipiuk 2003). Considering the Vangengeim-Girs hemispheric circulation indices for the European sector, seven circulation periods were distinguished (Degirmendžić et al. 2000).

The analyses of the distribution and variability of **atmospheric pressure** are the basis of research into atmospheric circulation. One of the first studies on the distribution of atmospheric pressure in Poland and Europe includes Gorczyński's work from (1917), in which the author presented 54 maps showing the distribution of monthly and annual isobars in Poland, Europe and all over the globe.

The Kraków series of measurements was subjected to the most detailed study of the meteorological element in question. It was used, among others, in Weisse's (1858), Hann's (1887), and Trepieńska's (1988) works. The oldest studies concerned mainly general comparisons of pressure values (their daily and annual courses as well as deviations from the average) between the various stations located in Poland and in Europe. Later studies, mainly Trepieńska's works also tackled the issues related

to the analysis of trends of changes since 1792 (Trepieńska 1988, 1997a; Barring et al. 2002). The author also made a number of comparisons of a long-term (since 1901) series of pressure measurements in Kraków and Warszawa, as well as in other European cities. Similar studies using a long series of measurements in Warszawa were also made by Ustrnul and Czekierda (2000). In the works mentioned above, it was found, among others, that the long-term variability of average annual pressure values shows a slight increase, which is mainly caused by pressure changes in the winter season. An important characteristic of the long-term variability of pressure throughout the year and in individual months is the occurrence of clear periods of higher or lower values, which is most likely associated with the occurrence of the so-called circular epochs.

Nevertheless, most studies using the atmospheric pressure measurements in Poland are based on shorter measurement series. They usually concern the impact of pressure and its changes on the human body (for example Koźmiński and Michalska 2012) or explain the changes and variability of other meteorological elements (for example Degirmendžić et al. 2004).

The **actinometric measurements** started in Poland at the end of the nineteenth century, but they were of experimental character and were short series. The oldest measurements of solar radiation were carried out in 1894 in the meteorological observatory in Puławy (Kolomijcov 1894). The pioneer of actinometric measurements in Poland, as well as the creator of the instruments used to measure the intensity of solar radiation (Moll-Gorczyński solarimeter, later known as Kipp & Zonen pyranometer), was professor Władysław Gorczyński. He initiated the long-term measurements of global solar radiation in Warszawa starting in 1900 (Gorczyński 1913), and then in Gdynia from 1920 (Gorczyński 1951).

It is known that short-term series of measurements were carried out in Grodzisk during the interwar period (Gorczyński 1911) in Sopot (1928–1935), Gdańsk (1931–1935) (Frischmuth 1935), in Kołobrzeg (Gorczyński 1951), Wrocław (1929–1932) (Grundmann 1933), Ursynów (Gorczyński 1951), Remiszewice near Łódź (June 1932), Januszewice near Opoczno (August–October 1932) (Gorczyński and Ostrowski 1934), Racibórz (1929–1941) (Mackiewicz 1957), the Sudetes Mountains (September–November 1931 and March–November 1932) (Stenz 1959) and in the Tatra Mountains (July–August 1903, January, April and September 1924 and 1935–1939) (Stenz 1925, 1959).

The measurements of solar radiation on a larger scale began in Poland only after the end of World War II, when in 1952, the State Hydrological and Meteorological Institute (PIHM) opened several actinometric stations (Bogdańska et al. 2002). However, it was difficult to carry out the analysis of the components of the radiation budget, especially its spatial distribution aspect, because the number of the stations was insufficient. Additionally, regular measurements were carried out in meteorological stations belonging to a few universities, the Polish Academy of Sciences Institutes (PAN) and the Institute of Soil Science and Plant Cultivation (IUNG). The development of actinometry in Poland, which was observed in the 1950s, provided the source material for numerous climatological research studies. These works were published on the basis of the research on long-term variability and regional differentiation of

solar radiation (Kuczmarska and Paszyński 1964a, b; Paszyński 1966; Podogrocki 1977, 1978; Miara et al. 1987; Olecki 1986, 1989; Słomka 1988; Bryś 1994) as well as the fluctuation, trends and periodicity analysis of global solar radiation (Bogdańska and Podogrocki 2000).

Until now, most of the **sunshine** studies have been based on short series of measurements, most often 10-year, and mainly includes the analysis of data from the 1960s and 1970s (Kuczmarski 1990) or earlier periods (Gorczyński 1913; Merecki 1914). Only a few stations in Poland can boast of long, homogeneous heliographic series. These include the IGiGP scientific station of the Jagiellonian University (formerly the Astronomical Observatory), where measurements have been taken continuously in the same place and using the same instrument since 1883. Based on the data from this station, numerous studies have been conducted on the subject of long-term variability of sunshine duration and its causes (among others, Morawska 1963; Matuszko 2014). In Kraków, as in other European cities (Matuszko 2016), one can notice common periods of decreases (“global dimming”: 1951–1980) and increases (“global brightening”: 1981–1995) in sunshine duration with extreme values occurring in the same years (maximum in 1943, 1921, minimum in 1980). Other stations with long, but partly reconstructed heliographic series, include Wrocław (Matuszko 2016), Śnieżka, Warszawa and Puławy (Bogdańska et al. 2002).

In addition, the various aspects of **cloudiness** variability in Poland have been analysed for many years, e.g. Satke (1898), Gorczyński and Wierzbicka (1915). Stenz (1952) and Okołowicz (1962) dealt with the problem of spatial variability of cloudiness amount in Poland. Warakomski (1969) reported on the seasonal course of cloud types in Poland. However, the number of papers dealing with the analysis of long-term changes in cloudiness has been very limited. Morawska-Horawska, in her analyses (1963, 1985), described in detail the observational series of cloudiness in Kraków, the longest in Poland, covering the period since 1859.

Air temperature is the element most often analysed and has one of the longest data series. In the case of this element, there have been countless analyses carried out on a local, regional and national (Polish) scale. The oldest and most important are studies by Gorczyński (1913, 1915a, b, 1916, 1918) and Gorczyński and Kosińska (1916), and in the later period by Romer (1947b, 1948/1949). Some of these studies are of particular importance because their results refer to spatial diversity across Europe. The issue of changes and variability of this element was considered for the numerous measuring stations that were located in areas currently in Poland (e.g. in Wrocław, Gdańsk, Poznań, Puławy, and on Śnieżka). However, most of the oldest studies presenting the variability of air temperature in different aspects concerned Warszawa and, especially, Kraków (Kowalczyk 1881; Karliński 1868, 1876; Kuczyński 1884; Merecki 1899). Nowadays, these series have been subjected to multiple detailed analyses. Studies for Kraków, edited by J. Trepińska (1997b), and those for Warszawa, the author of which is H. Lorenc (Lorenc 2000), contain their review. Owing to the complex history of measurements and observations, it is worth mentioning the temperature series for Gdansk (Miętus 2007; Filipiak 2007).

Air humidity is an element that plays an important role among meteorological processes within the atmosphere; however, the variety of humidity indices makes the

global view of air moisture changes difficult to ascertain. Moreover, some methodical difficulties in measuring and computing particular air humidity parameters have been often pointed out.

Despite the above-mentioned doubts about the reliability of the material at hand, studies on trends and distribution of air humidity parameters, in both regional and local aspects, have been conducted for many years. They present diurnal and annual trends in relative air humidity, saturation deficit and, rarely, in vapour pressure. The studies are most often excerpts from greater works, generally monographs of towns or regions. The data used in them are measurement series no longer than 30 years. Attempts to analyse the long-term variability of air humidity were based only on one of the parameters, usually the relative air humidity, or on much shorter observation series. The results describe tendencies at particular locations, and, up to the present, no research concerning the whole area of Poland has been undertaken.

The oldest works, in accordance with global trends, comprised mostly measurement and terminology problems (e.g. Gorczyński 1948; Demiańczuk 1963; Janiszewski 1975). Some studies on annual and seasonal distribution of humidity parameters were conducted on the regional or local scale (e.g. Wierzbicki 1878; Gumiński 1927; Kosiba 1952; Michna 1972), also regarding urban climate conditions (e.g. Tarajkowska 1974; Młostek and Sobik 1984; Gluza and Kaszewski 1984; Kłysik et al. 1985). Air humidity variability was seldom the topic of detailed examination in Poland. Hohendorf was the first (1967, 1969) to take up this topic and described changes in saturation deficit. B. Obrębska-Starkłowa and A. Grzyborowska (1997) presented the analysis of air humidity variability made for the plateau section of the Raba River (relative air humidity in the period of 1971–1992), T. Bryś (2003) described air saturation deficit variation in Wrocław over the twentieth century, and A. Wypych (e.g. 2004, 2010) focused on long-term variability of air humidity in Kraków, expressed by most of humidity parameters. Essentially, the conducted studies confirmed the drying of the atmosphere.

The oldest paper on **precipitation** variability in Poland by Kremser (1884) was based on chronological series starting in 1799 for Wrocław. Further research used the data from Warszawa (Wild 1887; Pietkiewicz 1889). Both authors found spatial differentiation in temporal variability of precipitation between 1813 and 1887. Analysing almost 100-year long precipitation series from Warszawa (1803–1910), Gorczyński (1911) established several precipitation epoch and mentioned the periodic character of precipitation fluctuations. The Warszawa series (1811–1910) was again analysed by Rychliński (1923a, b, 1924, 1927), who discussed indices describing precipitation variability year by year and its periodicity. Moreover, Rychliński (1927) found a diminishing degree of pluvial continentality between 1811 and 1910. This result was later proved by Romer (1947a) who found increased pluvial oceanity between 1851 and 1930 at some stations in Europe, including Poland.

The oldest works on **snow cover** in the Polish area originated at the turn of the nineteenth and twentieth centuries (Satke 1899; Kamińska 1912). They were, however, concerned mostly with the main characteristics of the snow cover and its physical properties, sometimes in individual winter seasons (Satke 1896; Kosińska-Bartnicka 1924). Research on the long-term variability of snow cover began to develop only

in the 1990s, with the progress of the investigations of the greenhouse effect and global warming. The first work on this subject was often published in the conference materials (e.g. Głowicki 1996; Piotrowicz 1996; Falarz et al. 1998), or even remained manuscripts (Głowicki and Jaskiewicz 1995). Detailed investigations of the long-term variability of different characteristics of the snow cover in Poznań for almost 70 winter seasons (1920/21-1989/90 with a war break) were conducted (Szustakowska 1991; Bednorz 1999a). As a result, a statistically significant positive trend of the period of potential snow cover duration in Poznań (3 days/10 years) was discovered, among others. It was launched the first research on the influence of atmospheric circulation on the snow cover long-term changes (Bednorz 1999b). A comparison of changes in snow conditions was conducted in the centre of a large city (Kraków) with the rural areas, resulting in a higher rate of decline in the values of snow-related characteristics in the city centre compared with rural areas (Falarz 1998). In addition, research was carried out in the mountainous areas of Poland (Głowicki 1996; Falarz et al. 1998). Within the 105-year snow cover series for the Śnieżka summit, a downward trend of the number of days with snow cover to the end of the 1930s, and a slight positive trend since the turn of the 1930s and 1940s has been observed (Głowicki and Jaskiewicz 1995). Polish-Slovak studies of snow cover on the southern and the northern slope of the Western Carpathians have shown a small negative trend in the number of days with snow cover depth ≥ 20 cm and in a maximum depth of snow in most of the investigated stations, except for the area of the Tatra mountains (Falarz et al. 1998). Research has been also done on changes and variability in snow conditions for skiing in the Polish Tatras (Falarz 1999).

Older climatological studies on spatial variability in the direction and speed of **wind** in Poland include the works of Stopa-Boryczka (1989), Niedzwiedz et al. (1985, 1995). At the end of the 1990s, research on climate change and variability in Poland began to emerge. However, in the climatological literature, studies describing the variability of temperature and precipitation in Poland dominate. Few items concern the issue of wind speed variability. Kożuchowski (2004) studied the variability of the speed of geostrophic wind and its components—meridian and zonal over Poland. Lorenc (1996, 2012) noted the increase in incidence of strong winds and whirlwinds. There are numerous publications concerning the forecasts of the change of wind field over Europe. However, the wind speed changes developed by the IPCC (2007) are not conclusive. Some models indicate an increase in the average and maximum wind speeds over Northern and Central Europe, others a decrease in wind. However, all forecasts developed by the IPCC highlight the possibility of a large seasonal wind speed variation, which is associated with a change in the pressure field over the Euro-Atlantic area.

The description of the occurrence of convective phenomena complements the climate characteristics of a given area. These phenomena include, above all, **thunderstorms**, accompanied by strong wind gusts and precipitation including hail, and tornadoes. Compared to most meteorological elements, the research of the phenomena mentioned above does not use measurement data, but mainly visual observations carried out at meteorological stations. Depending on the observation period, the records of such observations were of a different nature and were subject

to various changes. Until around the mid-twentieth century, descriptive characteristics could be encountered, although symbols were more and more commonly used in compliance with the rules for describing all atmospheric phenomena. Due to the difficulty in accessing detailed descriptions of the phenomena under consideration and the low level of importance attached to their relatively rare occurrence, the first more comprehensive study of the occurrence of thunderstorms and tornadoes in Poland appeared quite late. Initially, they were concerned about the occurrence of individual phenomena or very short observation periods (Smosarski 1915). The work of Wiszniewski (1949) is considered to be the first study in which longer observational series (1891–1930) was used, followed by the work of Stopa (1962, 1965) which concerns thunderstorms occurring in Poland in 1946–1955. These studies mainly took into account the spatial, annual and long-term diversity of thunderstorm occurrences. Later on, in the 1990s interest in thunderstorms increased again and the research was a continuation of the aforementioned works. However, while conducting these studies special attention was paid to the long-term (since 1885) changes in the occurrence of the described phenomenon (Bielec-Bąkowska 2003, 2013), to the synoptic conditions conducive to the formation of thunderstorms in Poland (Kolendowicz 2005) and to forecasts of their occurrence (Grabowska 2005). An important result of the research was the identification of thunderstorm regions in Poland and the demonstration of the existence of a tendency that the number of thunderstorm days varies depending on the region of the country. The majority of stations with a drop in the number of days with a storm are located northwest of the line connecting Śnieżka and Suwałki, while the remaining stations have positive trends. Recent advancements in technology have also facilitated instrumental monitoring of thunderstorm activity within the use of lightning detection systems. Thanks to PERUN network operating operationally in Poland since 2002, it was possible to develop high-resolution thunderstorm climatologies and explore previously undiscovered aspects such as storm intensity or diurnal cycles (Taszarek et al. 2015; Czernecki et al. 2016).

The occurrence of **tornadoes** in the area of Poland is the least investigated. Most of the works, including the earliest ones, were devoted to descriptions of individual cases (Gumiński 1936; Parczewski and Kluźniak 1959). More detailed studies, often covering longer periods of observation, were only started at the turn of the twenty-first century. They concerned both the spatial and temporal diversity of tornadoes (Lorenc 1996, 2012; Taszarek and Brooks 2015), environmental conditions in which they occur most often (Taszarek and Kolendowicz 2013) and case studies (Taszarek et al. 2016; Taszarek and Gromadzki 2017; Pilguy et al. 2019). Unfortunately, the short period from which reliable reports come from and spatial and temporal inhomogeneities do not allow to define reliable long-term trends in the frequency of tornadoes in Poland.

Overall, only a few studies have been devoted to the occurrence of storm precipitation, in particular **hail events**. The oldest studies include those devoted to hail tracks and areas of their most frequent occurrence (Schmuck 1959; Zinkiewicz and Michna 1955; Koźmiński and Rytel 1963). In later years, studies on hail and other storm precipitation types concerned mainly their long-term variability and diurnal cycles,

as well as the circulation conducive to their occurrence (Twardosz et al. 2011; Suwała 2014). On their basis, it was found that the most characteristic feature of storm precipitation is their annual course and the occurrence of clear fluctuations in long-term variability of the number of days and the thunderstorm precipitation totals (Twardosz et al. 2011; Bielec-Bąkowska 2014). However, apart from the southern regions of Poland, it is difficult to indicate clear trends in their frequency over longer periods of time (Twardosz et al. 2011; Bielec-Bąkowska 2014). In recent years, the application of machine learning techniques in radar data and numerical weather prediction models allowed for more detailed analysis of atmospheric potential for producing hail events in Poland (Czernecki et al. 2019). The rapid development of the European Severe Weather Database also enabled the assessment of large and very large hail events over the course of a few recent years (Pilorz 2015).

Lisowski and Bartnicki (1935) analysed the mean number of **days with fog**, for particular months, and for 19 localities. Their analyses covered additionally the frequency of fog in the periods spring-summer and autumn-winter in 28 stations placed in the areas which belonged to Poland before the Second World War. Gumiński (1952a) studied the annual course of fog occurrence in central and eastern Poland using data from the period 1929–1938. Prawdzic and Sucheta (1975) presented the mean number of days with fog in particular months and for the whole year, for 15 stations in the Pomorze and Mazury regions, for the period 1951–1960, while Piwkowski (1976) used data from 18 stations across the country and demonstrated the annual course of fog occurrence and the frequency of their origin and duration in seasons in the years 1961–1970. Morawska (1966) analysed data from the period 1861–1960 from Kraków and described multi-annual variability of the number of days with fog, together with the relationship between fog frequency, wind speed and air temperature. A similar study was published for Warsaw by Janiszewski (1967), with the application of data from the period 1948–1962.

Research of long-term changes of **bioclimatic conditions** is undertaken only in the last 20 years. Different biometeorological indices are used as the measures of bioclimate. Such studies were done by few research teams, mostly in the Institute of Geography and Spatial Organization of the Polish Academy of Sciences. As the first studies of the changes of bioclimatic conditions have based on data from Jagiellonian University station in Botanic Garden (UJBG) in Cracow. The data covered the period of 1901–2000. As the bioclimatic measures, the authors (Błażejczyk et al. 2003) have used Subjective Temperature (STI) and Insulation Predicted (Iclp) indices. The changes in STI and Iclp were analysed on the background of changes in air circulation. The research was continued by Błażejczyk and Twardosz (2010) who have studied changes in Wind Chill Temperature (*WCT*), *HUMIDEX*, accepted level of physical activity (*MHR*) as well as Physiological Subjective Temperature (*PST*) and Physiological Strain (*PhS*) indices at UJBG station in Cracow in the longer period of 1826–2006. Owczarek (2009) in her PhD thesis for the northern Poland has applied data for the period 1951–2000 to analyse changes in PhS, HL, STI, WCT and Heat Stress Index (HSI). In the last decade the changes of bioclimatic conditions were studied by Bąkowska (2011) for Kołobrzeg, Poznań and Szczawno